

表 124 2014年導入患者, 性別, 原疾患別

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	4,162	2,304	6,466		6,466
(%)	(16.9)	(19.5)	(17.8)		(17.8)
慢性腎盂腎炎	138	133	271		271
(%)	(0.6)	(1.1)	(0.7)		(0.7)
急速進行性糸球体腎炎	268	251	519		519
(%)	(1.1)	(2.1)	(1.4)		(1.4)
妊娠腎/妊娠中毒症		42	42		42
(%)		(0.4)	(0.1)		(0.1)
その他の分類不能の腎炎	82	64	146		146
(%)	(0.3)	(0.5)	(0.4)		(0.4)
多発性嚢胞腎	529	470	999		999
(%)	(2.2)	(4.0)	(2.7)		(2.7)
腎硬化症	3,542	1,609	5,151		5,151
(%)	(14.4)	(13.6)	(14.2)		(14.2)
悪性高血圧	205	110	315		315
(%)	(0.8)	(0.9)	(0.9)		(0.9)
糖尿病性腎症	11,424	4,385	15,809		15,809
(%)	(46.5)	(37.1)	(43.5)		(43.5)
SLE腎炎	98	170	268		268
(%)	(0.4)	(1.4)	(0.7)		(0.7)
アミロイド腎	39	49	88		88
(%)	(0.2)	(0.4)	(0.2)		(0.2)
痛風腎	67	6	73		73
(%)	(0.3)	(0.1)	(0.2)		(0.2)
先天性代謝異常による腎不全	9	14	23		23
(%)	(0.0)	(0.1)	(0.1)		(0.1)
腎・尿路結核	8	4	12		12
(%)	(0.0)	(0.0)	(0.0)		(0.0)
腎・尿路結石	37	26	63		63
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	138	40	178		178
(%)	(0.6)	(0.3)	(0.5)		(0.5)
閉塞性尿路障害	69	26	95		95
(%)	(0.3)	(0.2)	(0.3)		(0.3)
骨髄腫	76	56	132		132
(%)	(0.3)	(0.5)	(0.4)		(0.4)
腎形成不全	32	28	60		60
(%)	(0.1)	(0.2)	(0.2)		(0.2)
不明	2,612	1,490	4,102		4,102
(%)	(10.6)	(12.6)	(11.3)		(11.3)
再導入	130	74	204		204
(%)	(0.5)	(0.6)	(0.6)		(0.6)
その他	896	465	1,361		1,361
(%)	(3.6)	(3.9)	(3.7)		(3.7)
合計	24,561	11,816	36,377		36,377
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし					
総計	24,561	11,816	36,377		36,377

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査