

表 128 2014年導入患者, 原疾患別, 平均年齢

原疾患	人数	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	6,468	1	6,469	68.66	14.40
(%)	(17.8)	(7.7)	(17.8)		
慢性腎盂腎炎	270		270	68.86	14.60
(%)	(0.7)		(0.7)		
急速進行性糸球体腎炎	519		519	72.86	12.46
(%)	(1.4)		(1.4)		
妊娠腎／妊娠中毒症	42		42	64.81	14.13
(%)	(0.1)		(0.1)		
その他分類不能の腎炎	146		146	62.36	20.36
(%)	(0.4)		(0.4)		
多発性嚢胞腎	1,000		1,000	63.05	13.10
(%)	(2.7)		(2.7)		
腎硬化症	5,150		5,150	75.00	11.36
(%)	(14.2)		(14.2)		
悪性高血圧	315		315	64.95	16.72
(%)	(0.9)		(0.9)		
糖尿病性腎症	15,816		15,816	67.16	12.03
(%)	(43.5)		(43.5)		
SLE腎炎	268	1	269	63.47	16.44
(%)	(0.7)	(7.7)	(0.7)		
アミロイド腎	88		88	69.95	11.60
(%)	(0.2)		(0.2)		
痛風腎	72	1	73	68.43	10.75
(%)	(0.2)	(7.7)	(0.2)		
先天性代謝異常による腎不全	23		23	49.17	21.58
(%)	(0.1)		(0.1)		
腎・尿路結核	12		12	75.75	10.62
(%)	(0.0)		(0.0)		
腎・尿路結石	63		63	70.17	13.79
(%)	(0.2)		(0.2)		
腎・尿路腫瘍	178		178	72.91	10.90
(%)	(0.5)		(0.5)		
閉塞性尿路障害	95		95	70.07	13.71
(%)	(0.3)		(0.3)		
骨髄腫	132		132	71.60	11.73
(%)	(0.4)		(0.4)		
腎形成不全	60		60	39.82	24.64
(%)	(0.2)		(0.2)		
不明	4,094	10	4,104	72.03	13.53
(%)	(11.3)	(76.9)	(11.3)		
移植後再導入	208		208	57.61	16.20
(%)	(0.6)		(0.6)		
その他	1,363		1,363	69.30	15.27
(%)	(3.7)		(3.7)		
合計	36,382	13	36,395	69.04	13.39
(%)	(100.0)	(100.0)	(100.0)		
記載なし	0	10	10	0.00	0.00
総計	36,382	23	36,405	69.04	13.39

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査