

表 139 2014年導入患者死亡原因分類 (原疾患との関係)

原疾患	心不全	脳血管障害	感染症	消化管出血	悪性腫瘍	悪液質/尿毒症	心筋梗塞	カリウム中毒/頓死	肝硬変症	自殺/拒否	腸閉塞	肺血栓/肺塞栓	災害・事故死	その他	不明	合計	記載なし	総計
慢性糸球体腎炎	102	18	73	7	23	21	5	5	3	3	2	1	1	25	21	310		310
(%)	(32.9)	(5.8)	(23.5)	(2.3)	(7.4)	(6.8)	(1.6)	(1.6)	(1.0)	(1.0)	(0.6)	(0.3)	(0.3)	(8.1)	(6.8)	(100.0)		
慢性腎盂腎炎	2	1	7		3		4							5	1	23		23
(%)	(8.7)	(4.3)	(30.4)		(13.0)		(17.4)							(21.7)	(4.3)	(100.0)		
急速進行性糸球体腎炎	15	1	43	6	4	7	1		2	2				13	5	99		99
(%)	(15.2)	(1.0)	(43.4)	(6.1)	(4.0)	(7.1)	(1.0)		(2.0)	(2.0)				(13.1)	(5.1)	(100.0)		
妊娠腎/妊娠中毒症																		
(%)																		
その他の分類不能の腎炎	1	1	2	2										3	1	10		10
(%)	(10.0)	(10.0)	(20.0)	(20.0)										(30.0)	(10.0)	(100.0)		
多発性嚢胞腎	3	3	7		2	1								4	1	21		21
(%)	(14.3)	(14.3)	(33.3)		(9.5)	(4.8)								(19.0)	(4.8)	(100.0)		
腎硬化症	89	23	87	8	28	15	19	7	4	3	3	2	2	34	10	334		334
(%)	(26.6)	(6.9)	(26.0)	(2.4)	(8.4)	(4.5)	(5.7)	(2.1)	(1.2)	(0.9)	(0.9)	(0.6)	(0.6)	(10.2)	(3.0)	(100.0)		
悪性高血圧	2	1	1		1		1					1	1	1	3	12		12
(%)	(16.7)	(8.3)	(8.3)		(8.3)		(8.3)					(8.3)	(8.3)	(8.3)	(25.0)	(100.0)		
糖尿病性腎症	163	27	157	12	60	30	32	16	5	7	10	2	6	53	52	632		632
(%)	(25.8)	(4.3)	(24.8)	(1.9)	(9.5)	(4.7)	(5.1)	(2.5)	(0.8)	(1.1)	(1.6)	(0.3)	(0.9)	(8.4)	(8.2)	(100.0)		
SLE腎炎	6		11	1	1	3		1		1	1			2	3	30		30
(%)	(20.0)		(36.7)	(3.3)	(3.3)	(10.0)		(3.3)		(3.3)	(3.3)			(6.7)	(10.0)	(100.0)		
アミロイド腎	4		5	1		1		2					1	3		17		17
(%)	(23.5)		(29.4)	(5.9)		(5.9)		(11.8)					(5.9)	(17.6)		(100.0)		
痛風腎					1											1		1
(%)					(100.0)											(100.0)		
先天性代謝異常による腎不全																		
(%)																		
腎・尿路結核			1													1		1
(%)			(100.0)													(100.0)		
腎・尿路結石	2		2											1		5		5
(%)	(40.0)		(40.0)											(20.0)		(100.0)		
腎・尿路腫瘍	2	3	4		13	1				1	1			1		26		26
(%)	(7.7)	(11.5)	(15.4)		(50.0)	(3.8)				(3.8)	(3.8)			(3.8)		(100.0)		
閉塞性尿路障害	1	1	1		10	1						1			2	17		17
(%)	(5.9)	(5.9)	(5.9)		(58.8)	(5.9)						(5.9)			(11.8)	(100.0)		
骨髄腫	1	1	4		25									3	6	40		40
(%)	(2.5)	(2.5)	(10.0)		(82.5)									(7.5)	(15.0)	(100.0)		
腎形成不全	1		1											1	3	3		3
(%)	(33.3)		(33.3)											(33.3)		(100.0)		
不明	100	16	121	10	53	17	9	7	5	1	5	1		63	61	469		469
(%)	(21.3)	(3.4)	(25.8)	(2.1)	(11.3)	(3.6)	(1.9)	(1.5)	(1.1)	(0.2)	(1.1)	(0.2)		(13.4)	(13.0)	(100.0)		
再導入	5		2		2				1					2	1	13		13
(%)	(38.5)		(15.4)		(15.4)				(7.7)					(15.4)	(7.7)	(100.0)		
その他	48	8	63	5	37	2	4		5	3	3	1		44	13	236		236
(%)	(20.3)	(3.4)	(26.7)	(2.1)	(15.7)	(0.8)	(1.7)		(2.1)	(1.3)	(1.3)	(0.4)		(18.6)	(5.5)	(100.0)		
合計	547	104	592	52	263	99	75	38	25	21	25	9	11	258	180	2,299		2,299
(%)	(23.8)	(4.5)	(25.8)	(2.3)	(11.4)	(4.3)	(3.3)	(1.7)	(1.1)	(0.9)	(1.1)	(0.4)	(0.5)	(11.2)	(7.8)	(100.0)		
記載なし																		
(%)																		
総計	547	104	592	52	263	99	75	38	25	21	25	9	11	258	180	2,299		2,299
(%)	(23.8)	(4.5)	(25.8)	(2.3)	(11.4)	(4.3)	(3.3)	(1.7)	(1.1)	(0.9)	(1.1)	(0.4)	(0.5)	(11.2)	(7.8)	(100.0)		

数値下のかっこ内は行方向の合計に対する％です。
©Japanese Society for Dialysis Therapy
2014年調査