

表 145 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=多発性囊胞腎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)														1 (20.0)	2 (40.0)	1 (25.0)	2 (100.0)	2 (25.0)	1 (50.0)	3 (50.0)		3 (33.3)	3 (14.3)		3 (14.3)	81.33	10.02
脳血管障害 (%)														2 (100.0)	1 (100.0)	1 (50.0)	2 (20.0)	1 (20.0)	1 (25.0)	3 (50.0)		1 (33.3)	3 (14.3)		3 (14.3)	76.00	6.08
感染症 (%)																					7 (33.3)	7 (14.3)		7 (14.3)	78.43	11.27	
消化管出血 (%)																											
悪性腫瘍 (%)																											
悪液質/尿毒症 (%)																											
心筋梗塞 (%)																											
カリウム中毒/頓死 (%)																											
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																											
不明 (%)																											
合計 (%)										1 (100.0)	1 (100.0)		2 (100.0)	5 (100.0)	2 (100.0)	4 (100.0)	6 (100.0)		21 (100.0)				21 (100.0)	76.38	9.83		
記載なし																											
総計										1	1		2	5	2	4	6				21			21	76.38	9.83	

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査