

表 147 2014年導入患者死亡原因分類(年齢別, 原疾患別)原疾患=悪性高血圧

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																	1 (33.3)	1 (50.0)			2 (16.7)			2 (16.7)	83.50	2.12
脳血管障害 (%)																1 (50.0)					1 (8.3)			1 (8.3)	77.00	
感染症 (%)																	1 (33.3)				1 (8.3)			1 (8.3)	83.00	
消化管出血 (%)																										
悪性腫瘍 (%)													1 (100.0)								1 (8.3)			1 (8.3)	64.00	
悪液質/尿毒症 (%)																										
心筋梗塞 (%)																	1 (33.3)				1 (8.3)			1 (8.3)	84.00	
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																1 (50.0)					1 (8.3)			1 (8.3)	74.00	
災害・事故死 (%)												1 (50.0)									1 (8.3)			1 (8.3)	56.00	
その他 (%)																			1 (50.0)		1 (8.3)			1 (8.3)	86.00	
不明 (%)												1 (50.0)			1 (50.0)	1 (50.0)					3 (25.0)			3 (25.0)	68.33	9.29
合計 (%)												2 (100.0)	1 (100.0)		2 (100.0)	2 (100.0)	3 (100.0)	2 (100.0)			12 (100.0)			12 (100.0)	74.67	10.47
記載なし																										
総計												2	1		2	2	3	2			12			12	74.67	10.47

数値下のかっこ内は列方向の合計に対する%です。
 (C)Japanese Society for Dialysis Therapy
 2014年調査