

表 149 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=SLE腎炎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差		
心不全 (%)														2 (100.0)	1 (33.3)	1 (66.7)			1 (50.0)			5 (17.2)	1 (100.0)		6 (20.0)	75.20	10.76	
脳血管障害 (%)																												
感染症 (%)																									11 (36.7)	73.55	12.67	
消化管出血 (%)																									1 (3.3)	71.00		
悪性腫瘍 (%)																									1 (3.3)	57.00		
悪液質/尿毒症 (%)																									3 (10.0)	82.00	5.57	
心筋梗塞 (%)																												
カリウム中毒/頓死 (%)														1 (100.0)									1 (3.4)			1 (3.3)	40.00	
肝硬変症 (%)																												
自殺/拒否 (%)																									1 (3.3)	80.00		
腸閉塞 (%)																									1 (3.3)	76.00		
肺血栓/肺塞栓 (%)																												
災害・事故死 (%)																												
その他 (%)																									2 (6.7)	74.50	16.26	
不明 (%)																									3 (10.0)	73.67	14.74	
合計 (%)										1 (100.0)		1 (100.0)	3 (100.0)	3 (100.0)	2 (100.0)	3 (100.0)	6 (100.0)	4 (100.0)	4 (100.0)	2 (100.0)	29 (100.0)	1 (100.0)		30 (100.0)	73.28	12.69		
記載なし																												
総計										1		1	3	3	2	3	6	4	4	2		29	1		30	73.28	12.69	

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査