

表 155 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=腎・尿路腫瘍

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)															1 (16.7)	1 (50.0)	2 (7.7)				2	2	82.50	9.19		
脳血管障害 (%)														1 (20.0)	2 (33.3)					3 (11.5)	3 (11.5)	3	74.00	3.61		
感染症 (%)														2 (40.0)	1 (16.7)					4 (15.4)	4 (15.4)	4	78.75	8.02		
消化管出血 (%)																										
悪性腫瘍 (%)														1 (100.0)	3 (100.0)	2 (40.0)	2 (33.3)	2 (33.3)	1 (50.0)	1 (50.0)	1 (50.0)	13 (50.0)	13 (50.0)	13	73.00	11.49
悪液質/尿毒症 (%)																				1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	81.00	
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																				1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	84.00	
腸閉塞 (%)																				1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	83.00	
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																				1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	1 (3.8)	84.00	
不明 (%)																										
合計 (%)								1 (100.0)		1 (100.0)	3 (100.0)	5 (100.0)	6 (100.0)	6 (100.0)	2 (100.0)	2 (100.0)	2 (100.0)	2 (100.0)	26 (100.0)			26 (100.0)	26 (100.0)	26 (100.0)	76.27	9.66
記載なし																										
総計									1		1	3	5	6	6	2	2	2	26				26	76.27	9.66	

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査