

表 156 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=閉塞性尿路障害

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)																					1			1	88.00		
脳血管障害 (%)																					(5.9)			(5.9)			
感染症 (%)																					1			1	71.00		
消化管出血 (%)																					(5.9)			(5.9)			
悪性腫瘍 (%)																					1			1	66.00		
悪液質/尿毒症 (%)																					(5.9)			(5.9)			
心筋梗塞 (%)																					10			10	69.10	12.49	
カリウム中毒/頓死 (%)																					(58.8)			(58.8)			
肝硬変症 (%)																					1			1	85.00		
自殺/拒否 (%)																					(5.9)			(5.9)			
腸閉塞 (%)																					1			1	78.00		
肺血栓/肺塞栓 (%)																					(5.9)			(5.9)			
災害・事故死 (%)																					1			1	72.18	11.82	
その他 (%)																					2			2	74.00	15.56	
不明 (%)																					(11.8)			(11.8)			
合計 (%)										1 (100.0)			2 (100.0)	1 (100.0)	2 (100.0)	3 (100.0)	3 (100.0)	1 (100.0)	4 (100.0)			17 (100.0)			17	72.18	11.82
記載なし																											
総計										1			2	1	2	3	3	1	4			17			17	72.18	11.82

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査