

表 157 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=骨髄腫

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)																					1			1	81.00		
脳血管障害 (%)																					(2.5)			(2.5)			
感染症 (%)																					1			1	86.00		
消化管出血 (%)																					(2.5)			(2.5)			
悪性腫瘍 (%)																					4			4	65.75	31.40	
悪液質/尿毒症 (%)																					(10.0)			(10.0)			
カリウム中毒/頓死 (%)																					25			25	72.76	9.08	
心筋梗塞 (%)																					(62.5)			(62.5)			
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																					3			3	74.00	8.66	
不明 (%)																					(7.5)			(7.5)			
合計 (%)					1 (100.0)					2 (100.0)	1 (100.0)	4 (100.0)	7 (100.0)	6 (100.0)	10 (100.0)	5 (100.0)	3 (100.0)	1 (100.0)			40 (100.0)			40 (100.0)	71.67	12.36	
記載なし																											
総計					1					2	1	4	7	6	10	5	3	1			40			40	71.67	12.36	

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査