

表 158 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=腎形成不全

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差		
心不全 (%)																	1 (100.0)				1 (33.3)			1 (33.3)	82.00			
脳血管障害 (%)																									1 (33.3)	91.00		
感染症 (%)																									1 (33.3)			
消化管出血 (%)																												
悪性腫瘍 (%)																												
悪液質/尿毒症 (%)																												
カリウム中毒/頓死 (%)																												
心筋梗塞 (%)																												
肺血栓/肺塞栓 (%)																												
災害・事故死 (%)																												
その他 (%)																	1 (100.0)				1 (33.3)			1 (33.3)	74.00			
不明 (%)																												
合計 (%)																	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	3 (100.0)			3 (100.0)	82.33	8.50		
記載なし																												
総計																	1	1	1	1	3			3	82.33	8.50		

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査