

表 160 2014年導入患者死亡原因分類(年齢別、原疾患別)原疾患=移植後再導入

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)														2 (100.0)	2 (100.0)	1 (50.0)					5 (38.5)			5 (38.5)	73.40	6.35	
脳血管障害 (%)																							2 (15.4)	81.00	8.49		
感染症 (%)																1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)								
消化管出血 (%)																							2 (15.4)	52.50	6.36		
悪性腫瘍 (%)																							2 (15.4)	52.00			
悪液質/尿毒症 (%)																											
心筋梗塞 (%)																											
カリウム中毒/頓死 (%)																											
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																							2 (15.4)	86.00	5.66		
不明 (%)																							1 (7.7)	59.00			
合計 (%)										1 (100.0)	1 (100.0)	2 (100.0)		2 (100.0)	2 (100.0)	1 (100.0)	2 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	13 (100.0)			13 (100.0)	70.54	13.45
記載なし																											
総計										1	1	2		2	2	1	2	1	1	1	13			13	70.54	13.45	

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査