

表 162 2014年導入患者死亡原因分類(治療方法別)(2014年末)

死因	施設血液 透析	血液 透析 濾過	血液 濾過	血液 吸着 透析	在宅 血液 透析	腹膜透析	離 脱	生体 腎移植	献腎 移植	合 計	記 載 な し	総 計
心不全	203	4				5				212	333	545
(%)	(24.1)	(12.1)				(31.3)				(23.8)	(23.7)	(23.7)
脳血管障害	36	2								38	66	104
(%)	(4.3)	(6.1)								(4.3)	(4.7)	(4.5)
感染症	218	12		1		5				236	356	592
(%)	(25.9)	(36.4)		(100.0)		(31.3)				(26.5)	(25.4)	(25.8)
消化管出血	23					1				24	28	52
(%)	(2.7)					(6.3)				(2.7)	(2.0)	(2.3)
悪性腫瘍	95	3								98	164	262
(%)	(11.3)	(9.1)								(11.0)	(11.7)	(11.4)
悪液質/尿毒症	37	1								38	61	99
(%)	(4.4)	(3.0)								(4.3)	(4.3)	(4.3)
心筋梗塞	26					1				27	48	75
(%)	(3.1)					(6.3)				(3.0)	(3.4)	(3.3)
カリウム中毒/頓死	17	1				2				20	18	38
(%)	(2.0)	(3.0)				(12.5)				(2.2)	(1.3)	(1.7)
肝硬変症	7	1								8	17	25
(%)	(0.8)	(3.0)								(0.9)	(1.2)	(1.1)
自殺/拒否	8									8	13	21
(%)	(1.0)									(0.9)	(0.9)	(0.9)
腸閉塞	4	1								5	20	25
(%)	(0.5)	(3.0)								(0.6)	(1.4)	(1.1)
肺血栓/肺栓塞	3									3	6	9
(%)	(0.4)									(0.3)	(0.4)	(0.4)
災害死	6									6	5	11
(%)	(0.7)									(0.7)	(0.4)	(0.5)
その他	96	3								99	159	258
(%)	(11.4)	(9.1)								(11.1)	(11.3)	(11.2)
不明	63	5				2				70	110	180
(%)	(7.5)	(15.2)				(12.5)				(7.8)	(7.8)	(7.8)
合計	842	33		1		16				892	1,404	2,296
(%)	(100.0)	(100.0)		(100.0)		(100.0)				(100.0)	(100.0)	(100.0)
記載なし												
(%)												
総計	842	33		1		16				892	1,404	2,296
(%)												

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査