

表 212 2014年末患者, 性別, 原疾患別

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	58,148	38,822	96,970		96,970
(%)	(29.3)	(34.7)	(31.3)		(31.3)
慢性腎盂腎炎	1,398	1,644	3,042		3,042
(%)	(0.7)	(1.5)	(1.0)		(1.0)
急速進行性糸球体腎炎	1,279	1,215	2,494		2,494
(%)	(0.6)	(1.1)	(0.8)		(0.8)
妊娠腎/妊娠中毒症	11	1,603	1,614		1,614
(%)	(0.0)	(1.4)	(0.5)		(0.5)
その他の分類不能の腎炎	808	565	1,373		1,373
(%)	(0.4)	(0.5)	(0.4)		(0.4)
多発性嚢胞腎	5,985	5,021	11,006		11,006
(%)	(3.0)	(4.5)	(3.5)		(3.5)
腎硬化症	18,754	9,544	28,298		28,298
(%)	(9.5)	(8.5)	(9.1)		(9.1)
悪性高血圧	1,734	839	2,573		2,573
(%)	(0.9)	(0.7)	(0.8)		(0.8)
糖尿病性腎症	82,748	35,333	118,081		118,081
(%)	(41.8)	(31.6)	(38.1)		(38.1)
SLE腎炎	699	1,537	2,236		2,236
(%)	(0.4)	(1.4)	(0.7)		(0.7)
アミロイド腎	180	271	451		451
(%)	(0.1)	(0.2)	(0.1)		(0.1)
痛風腎	984	129	1,113		1,113
(%)	(0.5)	(0.1)	(0.4)		(0.4)
先天性代謝異常による腎不全	160	112	272		272
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結核	110	96	206		206
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結石	362	219	581		581
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	670	216	886		886
(%)	(0.3)	(0.2)	(0.3)		(0.3)
閉塞性尿路障害	471	252	723		723
(%)	(0.2)	(0.2)	(0.2)		(0.2)
骨髄腫	173	103	276		276
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎形成不全	401	248	649		649
(%)	(0.2)	(0.2)	(0.2)		(0.2)
不明	16,965	10,579	27,544		27,544
(%)	(8.6)	(9.4)	(8.9)		(8.9)
再導入	1,427	736	2,163		2,163
(%)	(0.7)	(0.7)	(0.7)		(0.7)
その他	4,674	2,883	7,557		7,557
(%)	(2.4)	(2.6)	(2.4)		(2.4)
合計	198,141	111,967	310,108		310,108
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし					
総計	198,141	111,967	310,108		310,108

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査