

表 219 2014年末患者，透析歴，原疾患別人数

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	12,803	16,492	21,217	16,426	12,189	8,024	4,782	3,049	1,630	350	96,962	8		96,970	11.10	9.23
(%)	(18.8)	(21.2)	(27.2)	(41.0)	(57.5)	(68.0)	(73.0)	(77.0)	(78.9)	(77.6)	(31.3)	(9.3)		(31.3)		
慢性腎盂腎炎	483	559	699	482	297	222	156	91	45	7	3,041	1		3,042	10.29	9.16
(%)	(0.7)	(0.7)	(0.9)	(1.2)	(1.4)	(1.9)	(2.4)	(2.3)	(2.2)	(1.6)	(1.0)	(1.2)		(1.0)		
急速進行性糸球体腎炎	734	721	563	247	112	65	28	15	6	3	2,494			2,494	5.60	6.36
(%)	(1.1)	(0.9)	(0.7)	(0.6)	(0.5)	(0.6)	(0.4)	(0.4)	(0.3)	(0.7)	(0.8)			(0.8)		
妊娠腎/妊娠中毒症	90	133	237	272	269	232	178	131	58	14	1,614			1,614	16.50	10.14
(%)	(0.1)	(0.2)	(0.3)	(0.7)	(1.3)	(2.0)	(2.7)	(3.3)	(2.8)	(3.1)	(0.5)			(0.5)		
その他の分類不能の腎炎	250	278	314	182	110	105	72	33	23	6	1,373			1,373	9.78	9.35
(%)	(0.4)	(0.4)	(0.4)	(0.5)	(0.5)	(0.9)	(1.1)	(0.8)	(1.1)	(1.3)	(0.4)			(0.4)		
多発性嚢胞腎	1,906	2,389	2,904	1,915	1,074	506	213	59	33	4	11,003	3		11,006	8.01	6.93
(%)	(2.8)	(3.1)	(3.7)	(4.8)	(5.1)	(4.3)	(3.3)	(1.5)	(1.6)	(0.9)	(3.5)	(3.5)		(3.5)		
腎硬化症	9,084	8,962	6,639	2,434	787	263	80	31	10	5	28,295	3		28,298	4.39	4.68
(%)	(13.3)	(11.5)	(8.5)	(6.1)	(3.7)	(2.2)	(1.2)	(0.8)	(0.5)	(1.1)	(9.1)	(3.5)		(9.1)		
悪性高血圧	565	671	649	338	199	89	43	10	4	2	2,570	3		2,573	6.78	6.61
(%)	(0.8)	(0.9)	(0.8)	(0.8)	(0.9)	(0.8)	(0.7)	(0.3)	(0.2)	(0.4)	(0.8)	(3.5)		(0.8)		
糖尿病性腎症	31,119	36,287	33,975	12,348	3,339	793	151	35	16	3	118,066	15		118,081	4.80	4.41
(%)	(45.7)	(46.6)	(43.6)	(30.8)	(15.7)	(6.7)	(2.3)	(0.9)	(0.8)	(0.7)	(38.1)	(17.4)		(38.1)		
SLE腎炎	456	458	504	340	223	125	76	38	14	1	2,235	1		2,236	8.65	8.09
(%)	(0.7)	(0.6)	(0.6)	(0.8)	(1.1)	(1.1)	(1.2)	(1.0)	(0.7)	(0.2)	(0.7)	(1.2)		(0.7)		
アミロイド腎	132	137	118	36	14	8	4	1	1		451			451	4.93	5.47
(%)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.0)	(0.0)		(0.1)			(0.1)		
痛風腎	165	215	267	198	151	62	37	11	4	3	1,113			1,113	9.35	7.72
(%)	(0.2)	(0.3)	(0.3)	(0.5)	(0.7)	(0.5)	(0.6)	(0.3)	(0.2)	(0.7)	(0.4)			(0.4)		
先天性代謝異常による腎不全	41	49	59	46	33	16	17	6	5		272			272	10.47	8.96
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	(0.3)	(0.2)	(0.2)		(0.1)			(0.1)		
腎・尿路結核	27	25	39	33	28	22	11	12	8	1	206			206	13.18	10.65
(%)	(0.0)	(0.0)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.3)	(0.4)	(0.2)	(0.1)			(0.1)		
腎・尿路結石	109	141	167	73	48	29	9	2	3		581			581	7.40	6.92
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)		(0.2)			(0.2)		
腎・尿路腫瘍	280	259	209	85	27	17	5	3	1		886			886	4.81	5.34
(%)	(0.4)	(0.3)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.0)		(0.3)			(0.3)		
閉塞性尿路障害	135	174	166	98	71	38	28	10	1	2	723			723	8.37	8.01
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.4)	(0.3)	(0.0)	(0.4)	(0.2)			(0.2)		
骨髄腫	152	78	32	11	1	1	1				276			276	2.40	3.56
(%)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)				(0.1)			(0.1)		
腎形成不全	102	111	124	94	94	54	38	20	10	2	649			649	11.16	9.45
(%)	(0.1)	(0.1)	(0.2)	(0.2)	(0.4)	(0.5)	(0.6)	(0.5)	(0.5)	(0.4)	(0.2)			(0.2)		
不明	7,081	7,342	6,777	3,257	1,586	729	354	229	109	30	27,494	50		27,544	6.27	6.71
(%)	(10.4)	(9.4)	(8.7)	(8.1)	(7.5)	(6.2)	(5.4)	(5.8)	(5.3)	(6.7)	(8.9)	(58.1)		(8.9)		
再導入	338	434	475	282	175	175	146	92	35	10	2,162	1		2,163	10.80	9.97
(%)	(0.5)	(0.6)	(0.6)	(0.7)	(0.8)	(1.5)	(2.2)	(2.3)	(1.7)	(2.2)	(0.7)	(1.2)		(0.7)		
その他	2,090	2,028	1,728	835	386	227	122	82	50	8	7,556	1		7,557	6.30	7.13
(%)	(3.1)	(2.6)	(2.2)	(2.1)	(1.8)	(1.9)	(1.9)	(2.1)	(2.4)	(1.8)	(2.4)	(1.2)		(2.4)		
合計	68,142	77,943	77,862	40,032	21,213	11,802	6,551	3,960	2,066	451	310,022	86		310,108	7.29	7.49
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)		
記載なし																
総計	68,142	77,943	77,862	40,032	21,213	11,802	6,551	3,960	2,066	451	310,022	86		310,108	7.29	7.49

数値下のかつこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査