

表 220 2014年末患者、透析歴、原疾患別人数(男性)

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	8,219	10,404	12,945	9,655	7,068	4,457	2,605	1,680	917	195	58,145	3		58,148	10.65	9.10
(%)	(17.8)	(19.9)	(25.9)	(39.7)	(58.0)	(70.0)	(75.5)	(80.3)	(81.9)	(81.3)	(29.4)	(4.9)		(29.3)		
慢性腎盂腎炎	237	275	328	200	129	96	59	42	27	4	1,397	1		1,398	9.90	9.25
(%)	(0.5)	(0.5)	(0.7)	(0.8)	(1.1)	(1.5)	(1.7)	(2.0)	(2.4)	(1.7)	(0.7)	(1.6)		(0.7)		
急速進行性糸球体腎炎	388	389	274	122	51	25	14	11	3	2	1,279			1,279	5.36	6.35
(%)	(0.8)	(0.7)	(0.5)	(0.5)	(0.4)	(0.4)	(0.4)	(0.5)	(0.3)	(0.8)	(0.6)			(0.6)		
妊娠腎/妊娠中毒症		5	1	2	1	1	1				11			11	9.45	8.95
(%)		(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)				(0.0)			(0.0)		
その他の分類不能の腎炎	145	166	173	110	70	64	40	20	17	3	808			808	9.99	9.48
(%)	(0.3)	(0.3)	(0.3)	(0.5)	(0.6)	(1.0)	(1.2)	(1.0)	(1.5)	(1.3)	(0.4)			(0.4)		
多発性嚢胞腎	1,066	1,358	1,562	1,036	531	262	111	30	24	3	5,983	2		5,985	7.82	6.90
(%)	(2.3)	(2.6)	(3.1)	(4.3)	(4.4)	(4.1)	(3.2)	(1.4)	(2.1)	(1.3)	(3.0)	(3.3)		(3.0)		
腎硬化症	6,170	6,026	4,249	1,598	489	145	49	20	4	2	18,752	2		18,754	4.26	4.57
(%)	(13.4)	(11.5)	(8.5)	(6.6)	(4.0)	(2.3)	(1.4)	(1.0)	(0.4)	(0.8)	(9.5)	(3.3)		(9.5)		
悪性高血圧	383	455	445	229	136	49	24	8	1	1	1,731	3		1,734	6.58	6.35
(%)	(0.8)	(0.9)	(0.9)	(0.9)	(1.1)	(0.8)	(0.7)	(0.4)	(0.1)	(0.4)	(0.9)	(4.9)		(0.9)		
糖尿病性腎症	22,448	25,994	23,457	8,188	2,086	452	87	14	9	1	82,736	12		82,748	4.63	4.27
(%)	(48.7)	(49.7)	(47.0)	(33.7)	(17.1)	(7.1)	(2.5)	(0.7)	(0.8)	(0.4)	(41.8)	(19.7)		(41.8)		
SLE腎炎	174	167	165	106	46	25	8	5	3		699			699	6.81	6.76
(%)	(0.4)	(0.3)	(0.3)	(0.4)	(0.4)	(0.4)	(0.2)	(0.2)	(0.3)		(0.4)			(0.4)		
アミロイド腎	59	59	39	8	8	1	4	1	1		180			180	4.77	6.23
(%)	(0.1)	(0.1)	(0.1)	(0.0)	(0.1)	(0.0)	(0.1)	(0.0)	(0.1)		(0.1)			(0.1)		
痛風腎	148	190	235	180	129	51	34	11	4	2	984			984	9.29	7.72
(%)	(0.3)	(0.4)	(0.5)	(0.7)	(1.1)	(0.8)	(1.0)	(0.5)	(0.4)	(0.8)	(0.5)			(0.5)		
先天性代謝異常による腎不全	20	32	29	30	21	10	11	4	3		160			160	10.97	9.04
(%)	(0.0)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.3)	(0.2)	(0.3)		(0.1)			(0.1)		
腎・尿路結核	21	9	19	21	15	11	6	5	2	1	110			110	12.42	10.22
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)	(0.4)	(0.1)			(0.1)		
腎・尿路結石	62	84	114	49	28	17	5	1	2		362			362	7.45	6.67
(%)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)	(0.1)	(0.0)	(0.2)		(0.2)			(0.2)		
腎・尿路腫瘍	215	211	150	60	17	11	3	2	1		670			670	4.55	5.14
(%)	(0.5)	(0.4)	(0.3)	(0.2)	(0.1)	(0.2)	(0.1)	(0.1)	(0.1)		(0.3)			(0.3)		
閉塞性尿路障害	98	118	110	62	37	22	16	7		1	471			471	7.73	7.77
(%)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.3)	(0.5)	(0.3)		(0.4)	(0.2)			(0.2)		
骨髄腫	98	53	13	7	1	1					173			173	2.18	3.29
(%)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)					(0.1)			(0.1)		
腎形成不全	61	61	92	58	55	34	22	14	4		401			401	11.01	9.08
(%)	(0.1)	(0.1)	(0.2)	(0.2)	(0.5)	(0.5)	(0.6)	(0.7)	(0.4)		(0.2)			(0.2)		
不明	4,520	4,666	4,151	1,934	915	392	183	107	46	15	16,929	36		16,965	5.92	6.35
(%)	(9.8)	(8.9)	(8.3)	(7.9)	(7.5)	(6.2)	(5.3)	(5.1)	(4.1)	(6.3)	(8.5)	(59.0)		(8.6)		
再導入	222	283	301	177	122	122	105	66	22	6	1,426	1		1,427	11.11	10.09
(%)	(0.5)	(0.5)	(0.6)	(0.7)	(1.0)	(1.9)	(3.0)	(3.2)	(2.0)	(2.5)	(0.7)	(1.6)		(0.7)		
その他	1,376	1,276	1,041	498	223	120	63	43	29	4	4,673	1		4,674	5.93	6.87
(%)	(3.0)	(2.4)	(2.1)	(2.0)	(1.8)	(1.9)	(1.8)	(2.1)	(2.6)	(1.7)	(2.4)	(1.6)		(2.4)		
合計	46,130	52,281	49,893	24,330	12,178	6,368	3,450	2,091	1,119	240	198,080	61		198,141	6.79	7.13
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)		
記載なし																
総計	46,130	52,281	49,893	24,330	12,178	6,368	3,450	2,091	1,119	240	198,080	61		198,141	6.79	7.13

数値下のかつこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査