

表 221 2014年末患者，透析歴，原疾患別人数(女性)

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	4,584	6,088	8,272	6,771	5,121	3,567	2,177	1,369	713	155	38,817	5		38,822	11.76	9.39
(%)	(20.8)	(23.7)	(29.6)	(43.1)	(56.7)	(65.6)	(70.2)	(73.2)	(75.3)	(73.5)	(34.7)	(20.0)		(34.7)		
慢性腎盂腎炎	246	284	371	282	168	126	97	49	18	3	1,644			1,644	10.61	9.07
(%)	(1.1)	(1.1)	(1.3)	(1.8)	(1.9)	(2.3)	(3.1)	(2.6)	(1.9)	(1.4)	(1.5)			(1.5)		
急速進行性糸球体腎炎	346	332	289	125	61	40	14	4	3	1	1,215			1,215	5.85	6.36
(%)	(1.6)	(1.3)	(1.0)	(0.8)	(0.7)	(0.7)	(0.5)	(0.2)	(0.3)	(0.5)	(1.1)			(1.1)		
妊娠腎/妊娠中毒症	90	128	236	270	268	231	177	131	58	14	1,603			1,603	16.55	10.14
(%)	(0.4)	(0.5)	(0.8)	(1.7)	(3.0)	(4.3)	(5.7)	(7.0)	(6.1)	(6.6)	(1.4)			(1.4)		
その他の分類不能の腎炎	105	112	141	72	40	41	32	13	6	3	565			565	9.48	9.18
(%)	(0.5)	(0.4)	(0.5)	(0.5)	(0.4)	(0.8)	(1.0)	(0.7)	(0.6)	(1.4)	(0.5)			(0.5)		
多発性嚢胞腎	840	1,031	1,342	879	543	244	102	29	9	1	5,020	1		5,021	8.24	6.96
(%)	(3.8)	(4.0)	(4.8)	(5.6)	(6.0)	(4.5)	(3.3)	(1.6)	(1.0)	(0.5)	(4.5)	(4.0)		(4.5)		
腎硬化症	2,914	2,936	2,390	836	298	118	31	11	6	3	9,543	1		9,544	4.65	4.89
(%)	(13.2)	(11.4)	(8.5)	(5.3)	(3.3)	(2.2)	(1.0)	(0.6)	(0.6)	(1.4)	(8.5)	(4.0)		(8.5)		
悪性高血圧	182	216	204	109	63	40	19	2	3	1	839			839	7.20	7.09
(%)	(0.8)	(0.8)	(0.7)	(0.7)	(0.7)	(0.7)	(0.6)	(0.1)	(0.3)	(0.5)	(0.7)			(0.7)		
糖尿病性腎症	8,671	10,293	10,518	4,160	1,253	341	64	21	7	2	35,330	3		35,333	5.19	4.70
(%)	(39.4)	(40.1)	(37.6)	(26.5)	(13.9)	(6.3)	(2.1)	(1.1)	(0.7)	(0.9)	(31.6)	(12.0)		(31.6)		
SLE腎炎	282	291	339	234	177	100	68	33	11	1	1,536	1		1,537	9.48	8.50
(%)	(1.3)	(1.1)	(1.2)	(1.5)	(2.0)	(1.8)	(2.2)	(1.8)	(1.2)	(0.5)	(1.4)	(4.0)		(1.4)		
アミロイド腎	73	78	79	28	6	7					271			271	5.03	4.90
(%)	(0.3)	(0.3)	(0.3)	(0.2)	(0.1)	(0.1)					(0.2)			(0.2)		
痛風腎	17	25	32	18	22	11	3			1	129			129	9.76	7.80
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.1)			(0.5)	(0.1)			(0.1)		
先天性代謝異常による腎不全	21	17	30	16	12	6	6	2	2		112			112	9.77	8.84
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	(0.2)		(0.1)			(0.1)		
腎・尿路結核	6	16	20	12	13	11	5	7	6		96			96	14.06	11.10
(%)	(0.0)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.4)	(0.6)		(0.1)			(0.1)		
腎・尿路結石	47	57	53	24	20	12	4	1	1		219			219	7.32	7.32
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)		(0.2)			(0.2)		
腎・尿路腫瘍	65	48	59	25	10	6	2	1			216			216	5.62	5.86
(%)	(0.3)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)			(0.2)			(0.2)		
閉塞性尿路障害	37	56	56	36	34	16	12	3	1	1	252			252	9.58	8.33
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.4)	(0.3)	(0.4)	(0.2)	(0.1)	(0.5)	(0.2)			(0.2)		
骨髄腫	54	25	19	4			1				103			103	2.78	3.95
(%)	(0.2)	(0.1)	(0.1)	(0.0)			(0.0)				(0.1)			(0.1)		
腎形成不全	41	50	32	36	39	20	16	6	6	2	248			248	11.40	10.03
(%)	(0.2)	(0.2)	(0.1)	(0.2)	(0.4)	(0.4)	(0.5)	(0.3)	(0.6)	(0.9)	(0.2)			(0.2)		
不明	2,561	2,676	2,626	1,323	671	337	171	122	63	15	10,565	14		10,579	6.83	7.23
(%)	(11.6)	(10.4)	(9.4)	(8.4)	(7.4)	(6.2)	(5.5)	(6.5)	(6.7)	(7.1)	(9.4)	(56.0)		(9.4)		
再導入	116	151	174	105	53	53	41	26	13	4	736			736	10.21	9.70
(%)	(0.5)	(0.6)	(0.6)	(0.7)	(0.6)	(1.0)	(1.3)	(1.4)	(1.4)	(1.9)	(0.7)			(0.7)		
その他	714	752	687	337	163	107	59	39	21	4	2,883			2,883	6.88	7.51
(%)	(3.2)	(2.9)	(2.5)	(2.1)	(1.8)	(2.0)	(1.9)	(2.1)	(2.2)	(1.9)	(2.6)			(2.6)		
合計	22,012	25,662	27,969	15,702	9,035	5,434	3,101	1,869	947	211	111,942	25		111,967	8.18	8.02
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)		
記載なし																
総計	22,012	25,662	27,969	15,702	9,035	5,434	3,101	1,869	947	211	111,942	25		111,967	8.18	8.02

数値下のかつこ内は列方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査