

表 226 2014年末患者、治療方法(新分類)、年齢別

治療方法 (新分類)	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
施設血液透析	3	1	13	62	176	512	1,331	3,084	6,503	10,150	13,844	19,853	32,113	42,526	41,396	36,452	28,238	14,865	3,937	564	255,623	18		255,641	68.17	12.35
(%)	(7.3)	(3.1)	(29.5)	(56.9)	(71.5)	(75.1)	(75.5)	(76.6)	(75.7)	(76.3)	(77.3)	(78.3)	(79.6)	(81.4)	(83.9)	(86.3)	(88.0)	(89.2)	(89.7)	(91.3)	(82.4)	(85.7)		(82.4)		
Off-line HDF				2	2	8	35	83	202	307	430	627	929	1,147	972	743	514	244	56	11	6,312	3		6,315	66.00	11.98
(%)			(1.8)	(0.8)	(1.2)	(2.0)	(2.1)	(2.3)	(2.3)	(2.4)	(2.5)	(2.3)	(2.2)	(2.0)	(1.8)	(1.6)	(1.5)	(1.3)	(1.8)	(2.0)	(14.3)			(2.0)		
On-line HDF	2			10	32	116	256	628	1,427	2,114	2,686	3,600	5,400	6,343	5,351	3,970	2,647	1,195	287	26	36,090			36,090	64.81	12.45
(%)	(4.9)			(9.2)	(13.0)	(17.0)	(14.5)	(15.6)	(16.6)	(15.9)	(15.0)	(14.2)	(13.4)	(12.1)	(10.8)	(9.4)	(8.2)	(7.2)	(6.5)	(4.2)	(11.6)			(11.6)		
Push/Pull HDF				1			2	7	12	21	25	39	65	106	91	71	58	32	4	3	537			537	68.23	11.93
(%)				(0.4)			(0.1)	(0.2)	(0.1)	(0.2)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(0.5)	(0.2)			(0.2)		
ハイオフィルトレーション				1			2	8	5	10	17	23	39	69	63	46	36	16	4	2	341			341	68.45	12.09
(%)				(0.4)			(0.1)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.3)	(0.1)			(0.1)		
血液濾過			1				1	1	1	4	7	5	10	20	12	15	9	2	3	1	92			92	67.63	13.69
(%)			(2.3)				(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.1)	(0.2)	(0.0)			(0.0)		
血液吸着透析 (リクセル)								2	2	24	72	149	341	458	338	169	60	6	2		1,623			1,623	66.63	7.53
(%)								(0.0)	(0.0)	(0.2)	(0.4)	(0.6)	(0.8)	(0.9)	(0.7)	(0.4)	(0.2)	(0.0)	(0.0)		(0.5)			(0.5)		
血液吸着透析 (その他)								1					4	2							7			7	60.00	9.80
(%)							(0.0)						(0.0)	(0.0)							(0.0)			(0.0)		
在宅血液透析				1	3	10	21	60	90	62	107	81	48	20	13	3	2				521			521	54.38	10.58
(%)				(0.4)	(0.4)	(0.6)	(0.5)	(0.7)	(0.7)	(0.3)	(0.4)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)			(0.2)			(0.2)		
腹膜透析 (CAPD)	1	1	3	7	8	20	56	99	194	303	392	497	748	914	699	475	350	188	60	6	5,021			5,021	64.29	13.04
(%)	(2.4)	(3.1)	(6.8)	(6.4)	(3.3)	(2.9)	(3.2)	(2.5)	(2.3)	(2.3)	(2.2)	(2.0)	(1.9)	(1.7)	(1.4)	(1.1)	(1.1)	(1.1)	(1.4)	(1.0)	(1.6)			(1.6)		
腹膜透析 (APD)	32	21	18	24	15	21	42	56	125	180	265	278	447	474	323	242	148	95	29	4	2,839			2,839	61.04	16.36
(%)	(78.0)	(65.6)	(40.9)	(22.0)	(6.1)	(3.1)	(2.4)	(1.4)	(1.5)	(1.4)	(1.5)	(1.1)	(1.1)	(0.9)	(0.7)	(0.6)	(0.5)	(0.6)	(0.7)	(0.6)	(0.9)			(0.9)		
腹膜透析 (CCPD)	3	9	9	4	10	2	29	35	65	102	113	172	175	152	102	55	24	14	5	1	1,081			1,081	57.58	14.48
(%)	(7.3)	(28.1)	(20.5)	(3.7)	(4.1)	(0.3)	(1.6)	(0.9)	(0.8)	(0.8)	(0.6)	(0.7)	(0.4)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.3)			(0.3)		
合計	41	32	44	109	246	682	1,764	4,025	8,596	13,305	17,913	25,350	40,352	52,259	49,367	42,251	32,087	16,659	4,387	618	310,087	21		310,108	67.54	12.49
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)			(100.0)		
記載なし																										
総計	41	32	44	109	246	682	1,764	4,025	8,596	13,305	17,913	25,350	40,352	52,259	49,367	42,251	32,087	16,659	4,387	618	310,087	21		310,108	67.54	12.49

数値下のかっこ内は列方向の合計に対する%です。

*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査