

表 241 2014年死亡患者(性別、原疾患)

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	4,384	2,758	7,142		7,142
(%)	(23.0)	(26.3)	(24.2)		(24.2)
慢性腎盂腎炎	118	125	243		243
(%)	(0.6)	(1.2)	(0.8)		(0.8)
急速進行性糸球体腎炎	186	154	340		340
(%)	(1.0)	(1.5)	(1.2)		(1.2)
妊娠腎/妊娠中毒症	3	65	68		68
(%)	(0.0)	(0.6)	(0.2)		(0.2)
その他の分類不能の腎炎	42	40	82		82
(%)	(0.2)	(0.4)	(0.3)		(0.3)
多発性嚢胞腎	321	260	581		581
(%)	(1.7)	(2.5)	(2.0)		(2.0)
腎硬化症	2,074	1,118	3,192		3,192
(%)	(10.9)	(10.7)	(10.8)		(10.8)
悪性高血圧	127	58	185		185
(%)	(0.7)	(0.6)	(0.6)		(0.6)
糖尿病性腎症	8,293	3,736	12,029		12,029
(%)	(43.6)	(35.6)	(40.7)		(40.7)
SLE腎炎	78	143	221		221
(%)	(0.4)	(1.4)	(0.7)		(0.7)
アミロイド腎	25	54	79		79
(%)	(0.1)	(0.5)	(0.3)		(0.3)
痛風腎	61	8	69		69
(%)	(0.3)	(0.1)	(0.2)		(0.2)
先天性代謝異常による腎不全	5	3	8		8
(%)	(0.0)	(0.0)	(0.0)		(0.0)
腎・尿路結核	13	15	28		28
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結石	39	21	60		60
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	116	29	145		145
(%)	(0.6)	(0.3)	(0.5)		(0.5)
閉塞性尿路障害	57	23	80		80
(%)	(0.3)	(0.2)	(0.3)		(0.3)
骨髄腫	71	40	111		111
(%)	(0.4)	(0.4)	(0.4)		(0.4)
腎形成不全	3	14	17		17
(%)	(0.0)	(0.1)	(0.1)		(0.1)
不明	2,321	1,419	3,740		3,740
(%)	(12.2)	(13.5)	(12.7)		(12.7)
再導入	70	32	102		102
(%)	(0.4)	(0.3)	(0.3)		(0.3)
その他	622	381	1,003		1,003
(%)	(3.3)	(3.6)	(3.4)		(3.4)
合計	19,029	10,496	29,525		29,525
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし					
総計	19,029	10,496	29,525		29,525

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査