

表 249 2014年死亡患者(男性 透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1 年 ～	2 年 ～	5 年 ～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	114	93	159	312	833	1,027	669		691	467	4,365	2	17	4,384
(%)	(12.3)	(13.1)	(15.2)	(16.9)	(17.6)	(19.7)	(28.2)		(46.6)	(72.7)	(23.0)	(9.1)	(33.3)	(23.0)
慢性腎盂腎炎	8	4	11	11	21	27	10		23	3	118			118
(%)	(0.9)	(0.6)	(1.1)	(0.6)	(0.4)	(0.5)	(0.4)		(1.6)	(0.5)	(0.6)			(0.6)
急速進行性糸球体腎炎	49	15	11	18	47	32	7		3	4	186			186
(%)	(5.3)	(2.1)	(1.1)	(1.0)	(1.0)	(0.6)	(0.3)		(0.2)	(0.6)	(1.0)			(1.0)
妊娠腎/妊娠中毒症									1	2	3			3
(%)									(0.1)	(0.3)	(0.0)			(0.0)
その他の分類不能の腎炎	4		4		10	9	7		5	2	41		1	42
(%)	(0.4)		(0.4)		(0.2)	(0.2)	(0.3)		(0.3)	(0.3)	(0.2)		(2.0)	(0.2)
多発性嚢胞腎	9	8	8	16	69	65	57		65	23	320		1	321
(%)	(1.0)	(1.1)	(0.8)	(0.9)	(1.5)	(1.2)	(2.4)		(4.4)	(3.6)	(1.7)		(2.0)	(1.7)
腎硬化症	124	126	164	257	638	505	176		73	7	2,070		4	2,074
(%)	(13.4)	(17.8)	(15.7)	(14.0)	(13.5)	(9.7)	(7.4)		(4.9)	(1.1)	(10.9)		(7.8)	(10.9)
悪性高血圧	7		3	15	30	35	18		16	3	127			127
(%)	(0.8)		(0.3)	(0.8)	(0.6)	(0.7)	(0.8)		(1.1)	(0.5)	(0.7)			(0.7)
糖尿病性腎症	247	247	394	842	2,278	2,758	1,092		396	22	8,276	3	14	8,293
(%)	(26.6)	(34.8)	(37.8)	(45.7)	(48.2)	(52.9)	(46.0)		(26.7)	(3.4)	(43.7)	(13.6)	(27.5)	(43.6)
SLE腎炎	8	5	7	5	18	21	6		7	1	78			78
(%)	(0.9)	(0.7)	(0.7)	(0.3)	(0.4)	(0.4)	(0.3)		(0.5)	(0.2)	(0.4)			(0.4)
アミロイド腎	1	4	3	2	8	5	2				25			25
(%)	(0.1)	(0.6)	(0.3)	(0.1)	(0.2)	(0.1)	(0.1)				(0.1)			(0.1)
痛風腎	2		1	2	8	15	12		14	6	60	1		61
(%)	(0.2)		(0.1)	(0.1)	(0.2)	(0.3)	(0.5)		(0.9)	(0.9)	(0.3)	(4.5)		(0.3)
先天性代謝異常による腎不全					1	1	2			1	5			5
(%)					(0.0)	(0.0)	(0.1)			(0.2)	(0.0)			(0.0)
腎・尿路結核			1	1	4	2	2		2	1	13			13
(%)			(0.1)	(0.1)	(0.1)	(0.0)	(0.1)		(0.1)	(0.2)	(0.1)			(0.1)
腎・尿路結石	2	2	1	4	7	12	6		4	1	39			39
(%)	(0.2)	(0.3)	(0.1)	(0.2)	(0.1)	(0.2)	(0.3)		(0.3)	(0.2)	(0.2)			(0.2)
腎・尿路腫瘍	17	9	14	16	33	13	7		5		114	1	1	116
(%)	(1.8)	(1.3)	(1.3)	(0.9)	(0.7)	(0.2)	(0.3)		(0.3)		(0.6)	(4.5)	(2.0)	(0.6)
閉塞性尿路障害	10	3	3	5	7	20	6		3		57			57
(%)	(1.1)	(0.4)	(0.3)	(0.3)	(0.1)	(0.4)	(0.3)		(0.2)		(0.3)			(0.3)
骨髄腫	17	10	16	13	8	5	1				70		1	71
(%)	(1.8)	(1.4)	(1.5)	(0.7)	(0.2)	(0.1)	(0.0)				(0.4)		(2.0)	(0.4)
腎形成不全					2					1	3			3
(%)					(0.0)					(0.2)	(0.0)			(0.0)
不明	192	121	174	249	564	540	244		135	79	2,298	13	10	2,321
(%)	(20.7)	(17.1)	(16.7)	(13.5)	(11.9)	(10.4)	(10.3)		(9.1)	(12.3)	(12.1)	(59.1)	(19.6)	(12.2)
再導入	6	6	5	6	11	15	6		10	5	70			70
(%)	(0.6)	(0.8)	(0.5)	(0.3)	(0.2)	(0.3)	(0.3)		(0.7)	(0.8)	(0.4)			(0.4)
その他	110	56	64	68	128	103	46		29	14	618	2	2	622
(%)	(11.9)	(7.9)	(6.1)	(3.7)	(2.7)	(2.0)	(1.9)		(2.0)	(2.2)	(3.3)	(9.1)	(3.9)	(3.3)
合計	927	709	1,043	1,842	4,725	5,210	2,376		1,482	642	18,956	22	51	19,029
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし														
総計	927	709	1,043	1,842	4,725	5,210	2,376		1,482	642	18,956	22	51	19,029

数値下のかっこ内は列方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査