

表 250 2014年死亡患者(女性 透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1 年 ～	2 年 ～	5 年 ～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	90	51	87	166	446	643	438		499	331	2,751	1	6	2,758
(%)	(15.5)	(13.6)	(17.5)	(20.2)	(19.5)	(22.0)	(29.9)		(48.3)	(66.6)	(26.3)	(25.0)	(30.0)	(26.3)
慢性腎盂腎炎	9	6	7	7	17	25	20		23	10	124		1	125
(%)	(1.5)	(1.6)	(1.4)	(0.9)	(0.7)	(0.9)	(1.4)		(2.2)	(2.0)	(1.2)		(5.0)	(1.2)
急速進行性糸球体腎炎	35	13	13	17	27	34	8		7		154			154
(%)	(6.0)	(3.5)	(2.6)	(2.1)	(1.2)	(1.2)	(0.5)		(0.7)		(1.5)			(1.5)
妊娠腎/妊娠中毒症			1	1	4	7	9		15	27	64		1	65
(%)			(0.2)	(0.1)	(0.2)	(0.2)	(0.6)		(1.5)	(5.4)	(0.6)		(5.0)	(0.6)
その他の分類不能の腎炎	6	3	2	3	5	11	5		2	3	40			40
(%)	(1.0)	(0.8)	(0.4)	(0.4)	(0.2)	(0.4)	(0.3)		(0.2)	(0.6)	(0.4)			(0.4)
多発性嚢胞腎	5	2	13	12	42	65	53		49	19	260			260
(%)	(0.9)	(0.5)	(2.6)	(1.5)	(1.8)	(2.2)	(3.6)		(4.7)	(3.8)	(2.5)			(2.5)
腎硬化症	56	49	73	117	339	303	122		47	10	1,116		2	1,118
(%)	(9.6)	(13.0)	(14.7)	(14.3)	(14.8)	(10.4)	(8.3)		(4.5)	(2.0)	(10.7)		(10.0)	(10.7)
悪性高血圧	3	2	7	5	13	13	10		4	1	58			58
(%)	(0.5)	(0.5)	(1.4)	(0.6)	(0.6)	(0.4)	(0.7)		(0.4)	(0.2)	(0.6)			(0.6)
糖尿病性腎症	112	100	151	301	926	1,314	586		216	20	3,726	1	9	3,736
(%)	(19.3)	(26.6)	(30.4)	(36.7)	(40.5)	(45.0)	(40.1)		(20.9)	(4.0)	(35.6)	(25.0)	(45.0)	(35.6)
SLE腎炎	15	4	8	7	26	40	13		21	9	143			143
(%)	(2.6)	(1.1)	(1.6)	(0.9)	(1.1)	(1.4)	(0.9)		(2.0)	(1.8)	(1.4)			(1.4)
アミロイド腎	8	4	9	5	10	12	3		3		54			54
(%)	(1.4)	(1.1)	(1.8)	(0.6)	(0.4)	(0.4)	(0.2)		(0.3)		(0.5)			(0.5)
痛風腎				1	3	1	1		1	1	8			8
(%)				(0.1)	(0.1)	(0.0)	(0.1)		(0.1)	(0.2)	(0.1)			(0.1)
先天性代謝異常による腎不全					2				1		3			3
(%)					(0.1)				(0.1)		(0.0)			(0.0)
腎・尿路結核	1		1			2	3		5	3	15			15
(%)	(0.2)		(0.2)			(0.1)	(0.2)		(0.5)	(0.6)	(0.1)			(0.1)
腎・尿路結石		1	1	3	7	4	2		1	2	21			21
(%)		(0.3)	(0.2)	(0.4)	(0.3)	(0.1)	(0.1)		(0.1)	(0.4)	(0.2)			(0.2)
腎・尿路腫瘍	2	2	2		11	6	3		2	1	29			29
(%)	(0.3)	(0.5)	(0.4)		(0.5)	(0.2)	(0.2)		(0.2)	(0.2)	(0.3)			(0.3)
閉塞性尿路障害	4	2		2	5	3	3		2	2	23			23
(%)	(0.7)	(0.5)		(0.2)	(0.2)	(0.1)	(0.2)		(0.2)	(0.4)	(0.2)			(0.2)
骨髄腫	7	14	7	1	7	3	1				40			40
(%)	(1.2)	(3.7)	(1.4)	(0.1)	(0.3)	(0.1)	(0.1)				(0.4)			(0.4)
腎形成不全	1	2	1		1	1	5		1	2	14			14
(%)	(0.2)	(0.5)	(0.2)		(0.0)	(0.0)	(0.3)		(0.1)	(0.4)	(0.1)			(0.1)
不明	152	77	80	133	321	362	142		105	44	1,416	2	1	1,419
(%)	(26.2)	(20.5)	(16.1)	(16.2)	(14.0)	(12.4)	(9.7)		(10.2)	(8.9)	(13.5)	(50.0)	(5.0)	(13.5)
再導入	5	2	1	5	8	7				4	32			32
(%)	(0.9)	(0.5)	(0.2)	(0.6)	(0.4)	(0.2)				(0.8)	(0.3)			(0.3)
その他	70	42	33	35	65	63	36		29	8	381			381
(%)	(12.0)	(11.2)	(6.6)	(4.3)	(2.8)	(2.2)	(2.5)		(2.8)	(1.6)	(3.6)			(3.6)
合計	581	376	497	821	2,285	2,919	1,463		1,033	497	10,472	4	20	10,496
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし														
総計	581	376	497	821	2,285	2,919	1,463		1,033	497	10,472	4	20	10,496

数値下のかっこ内は列方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査