

表 301 2014年死亡患者死亡原因分類（原疾患との関係）

原疾患	心不全	脳血管障害	感染症	消化管出血	悪性腫瘍	悪液質/尿毒症	心筋梗塞	カリウム中毒/頓死	肝硬変症	自殺/拒否	腸閉塞	肺血栓/肺塞栓	災害・事故死	その他	不明	合計	記載なし	総計
慢性糸球体腎炎	2,089	522	1,414	105	612	294	216	151	83	53	87	19	42	666	789	7,142		7,142
(%)	(29.2)	(7.3)	(19.8)	(1.5)	(8.6)	(4.1)	(3.0)	(2.1)	(1.2)	(0.7)	(1.2)	(0.3)	(0.6)	(9.3)	(11.0)	(100.0)		(100.0)
慢性腎盂腎炎	45	21	53	4	24	5	13	4	1	1	2	1	2	34	33	243		243
(%)	(18.5)	(8.6)	(21.8)	(1.6)	(9.9)	(2.1)	(5.3)	(1.6)	(0.4)	(0.4)	(0.8)	(0.4)	(0.8)	(14.0)	(13.6)	(100.0)		(100.0)
急速進行性糸球体腎炎	66	23	110	8	24	17	5	11	4	5	9		2	39	17	340		340
(%)	(19.4)	(8.8)	(32.4)	(2.4)	(7.1)	(5.0)	(1.5)	(3.2)	(1.2)	(1.5)	(2.6)		(0.6)	(11.5)	(5.0)	(100.0)		(100.0)
妊娠腎/妊娠中毒症	25	4	16		4		4	3						4	8	68		68
(%)	(36.8)	(5.9)	(23.5)		(5.9)		(5.9)	(4.4)						(5.9)	(11.8)	(100.0)		(100.0)
その他の分類不能の腎炎	13	7	18	4	7	3	4	3		2				12	9	82		82
(%)	(15.9)	(8.5)	(22.0)	(4.9)	(8.5)	(3.7)	(4.9)	(3.7)		(2.4)				(14.6)	(11.0)	(100.0)		(100.0)
多発性囊胞腎	128	56	140	15	41	21	15	10	6	4	5	2	1	71	66	581		581
(%)	(22.0)	(9.6)	(24.1)	(2.6)	(7.1)	(3.8)	(2.6)	(1.7)	(1.0)	(0.7)	(0.9)	(0.3)	(0.2)	(12.2)	(11.4)	(100.0)		(100.0)
腎硬化症	848	197	689	48	313	146	133	82	24	24	31	9	22	341	285	3,192		3,192
(%)	(26.6)	(6.2)	(21.6)	(1.5)	(9.8)	(4.6)	(4.2)	(2.6)	(0.8)	(0.8)	(1.0)	(0.3)	(0.7)	(10.7)	(8.9)	(100.0)		(100.0)
悪性高血圧	37	16	32	2	18	9	6	7	2		1	3	2	20	30	185		185
(%)	(20.0)	(8.6)	(17.3)	(1.1)	(9.7)	(4.9)	(3.2)	(3.8)	(1.1)		(0.5)	(1.6)	(1.1)	(10.8)	(16.2)	(100.0)		(100.0)
糖尿病性腎症	3,243	907	2,492	169	959	464	718	400	103	93	107	30	58	930	1,356	12,029		12,029
(%)	(27.0)	(7.5)	(20.7)	(1.4)	(8.0)	(3.9)	(6.0)	(3.3)	(0.9)	(0.8)	(0.9)	(0.2)	(0.5)	(7.7)	(11.3)	(100.0)		(100.0)
SLC腎炎	54	7	58	7	13	11	9	7		3	3			26	23	221		221
(%)	(24.4)	(3.2)	(26.2)	(3.2)	(5.9)	(5.0)	(4.1)	(3.2)		(1.4)	(1.4)			(11.8)	(10.4)	(100.0)		(100.0)
アミロイド腎	24	1	17	3	3	1	1	6				1		1	14	6	79	79
(%)	(30.4)	(1.3)	(21.5)	(3.8)	(3.8)	(1.3)	(1.3)	(7.6)			(1.3)	(1.3)	(1.3)	(17.7)	(7.6)	(100.0)		(100.0)
痛風腎	16	10	9	1	10	5		2	1			1		8	6	69		69
(%)	(23.2)	(14.5)	(13.0)	(1.4)	(14.5)	(7.2)		(2.9)	(1.4)			(1.4)		(11.6)	(8.7)	(100.0)		(100.0)
先天性代謝異常による腎不全	2		1		2			2							1	8		8
(%)	(25.0)		(12.5)		(25.0)			(25.0)							(12.5)	(100.0)		(100.0)
腎・尿路結核	6		6	2	2	1	1	1					1	4	4	28		28
(%)	(21.4)		(21.4)	(7.1)	(7.1)	(3.6)	(3.6)	(3.6)					(3.6)	(14.3)	(14.3)	(100.0)		(100.0)
腎・尿路結石	13	4	14	1	6	4	1		1	2				5	9	60		60
(%)	(21.7)	(6.7)	(23.3)	(1.7)	(10.0)	(6.7)	(1.7)		(1.7)	(3.3)				(8.3)	(15.0)	(100.0)		(100.0)
腎・尿路腫瘍	22	8	17	1	61	6	2	4	1	1	2			11	9	145		145
(%)	(15.2)	(5.5)	(11.7)	(0.7)	(42.1)	(4.1)	(1.4)	(2.8)	(0.7)	(0.7)	(1.4)			(7.6)	(6.2)	(100.0)		(100.0)
閉塞性尿路障害	18	6	12	2	16	3	4	1				1		6	11	80		80
(%)	(22.5)	(7.5)	(15.0)	(2.5)	(20.0)	(3.8)	(5.0)	(1.3)				(1.3)		(7.5)	(13.8)	(100.0)		(100.0)
骨髄腫	4	4	11		70	1								6	14	111		111
(%)	(3.6)	(3.6)	(9.9)		(63.1)	(0.9)		(0.9)						(5.4)	(12.6)	(100.0)		(100.0)
腎形成不全	2	3	5		2				1		1			3		17		17
(%)	(11.8)	(17.6)	(29.4)		(11.8)				(5.9)		(5.9)			(17.6)		(100.0)		(100.0)
不明	896	239	805	75	336	147	92	81	31	16	47	8	23	400	544	3,740		3,740
(%)	(24.0)	(8.4)	(21.5)	(2.0)	(9.0)	(3.9)	(2.5)	(2.2)	(0.8)	(0.4)	(1.3)	(0.2)	(0.6)	(10.7)	(14.5)	(100.0)		(100.0)
再導入	24	7	22	1	8	3	5	5	3	2	1			11	10	102		102
(%)	(23.5)	(6.9)	(21.6)	(1.0)	(7.8)	(2.9)	(4.9)	(4.9)	(2.9)	(2.0)	(1.0)			(10.8)	(9.8)	(100.0)		(100.0)
その他	200	56	221	23	138	29	29	13	17	9	13	4	3	139	109	1,003		1,003
(%)	(19.9)	(5.8)	(22.0)	(2.3)	(13.8)	(2.9)	(2.9)	(1.3)	(1.7)	(0.9)	(1.3)	(0.4)	(0.3)	(13.9)	(10.9)	(100.0)		(100.0)
合計	7,775	2,098	6,162	471	2,669	1,170	1,258	794	278	215	310	79	157	2,750	3,339	29,525		29,525
(%)	(26.3)	(7.1)	(20.9)	(1.6)	(9.9)	(4.0)	(4.3)	(2.7)	(0.9)	(0.7)	(1.0)	(0.3)	(0.5)	(9.3)	(11.3)	(100.0)		(100.0)
記載なし																		
総計	7,775	2,098	6,162	471	2,669	1,170	1,258	794	278	215	310	79	157	2,750	3,339	29,525		29,525
(%)	(26.3)	(7.1)	(20.9)	(1.6)	(9.9)	(4.0)	(4.3)	(2.7)	(0.9)	(0.7)	(1.0)	(0.3)	(0.5)	(9.3)	(11.3)	(100.0)		(100.0)

数値下のかっこ内は行方向の合計に対する％です。
©Japanese Society for Dialysis Therapy
2014年調査