

表 305 2014年死亡患者死亡原因分類(年齢別、原疾患別)原疾患=妊娠腎／妊娠中毒症

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)											3 (50.0)	4 (40.0)	3 (33.3)	5 (26.3)	4 (44.4)	5 (50.0)	1 (33.3)				25 (36.8)			25 (36.8)	71.24	8.49	
脳血管障害 (%)											1 (10.0)		3 (15.8)								4 (5.9)			4 (5.9)	70.75	5.19	
感染症 (%)											1 (16.7)	4 (40.0)	2 (22.2)	4 (21.1)		4 (40.0)	1 (33.3)				16 (23.5)			16 (23.5)	71.69	8.91	
消化管出血 (%)											1 (100.0)	1 (16.7)															
悪性腫瘍 (%)																2 (22.2)					4 (5.9)			4 (5.9)	66.00	13.39	
悪液質/尿毒症 (%)																											
心筋梗塞 (%)																					4 (5.9)			4 (5.9)	70.25	1.71	
カリウム中毒/頓死 (%)																					3 (4.4)			3 (4.4)	72.00	2.65	
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)											1 (16.7)	1 (10.0)	1 (11.1)	1 (5.3)								4 (5.9)			4 (5.9)	64.50	4.65
不明 (%)											1 (100.0)		2 (22.2)	1 (5.3)	2 (22.2)	1 (10.0)	1 (33.3)				8 (11.8)			8 (11.8)	72.38	11.58	
合計 (%)											1 (100.0)	1 (100.0)	6 (100.0)	10 (100.0)	9 (100.0)	19 (100.0)	9 (100.0)	10 (100.0)	3 (100.0)			68 (100.0)			68 (100.0)	70.72	8.52
記載なし																											
総計											1	1	6	10	9	19	9	10	3			68			68	70.72	8.52

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査