

表 312 2014年死亡患者死亡原因分類(年齢別, 原疾患別)原疾患＝アミロイド腎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)												1 (33.3)	3 (27.3)	3 (23.1)	6 (50.0)	8 (42.1)	2 (13.3)	1 (50.0)			24 (30.4)			24 (30.4)	72.63	7.33
脳血管障害 (%)												1 (33.3)									1 (1.3)			1 (1.3)	59.00	
感染症 (%)													2 (18.2)	4 (30.8)	1 (8.3)	5 (26.3)	5 (33.3)				17 (21.5)			17 (21.5)	73.94	7.12
消化管出血 (%)									1 (100.0)				2 (18.2)								3 (3.8)			3 (3.8)	55.67	10.12
悪性腫瘍 (%)								1 (100.0)										2 (13.3)			3 (3.8)			3 (3.8)	67.33	26.31
悪液質/尿毒症 (%)																1 (5.3)					1 (1.3)			1 (1.3)	77.00	
心筋梗塞 (%)															1 (8.3)						1 (1.3)			1 (1.3)	71.00	
カリウム中毒/頓死 (%)													1 (9.1)	3 (23.1)		1 (5.3)	1 (6.7)				6 (7.6)			6 (7.6)	69.83	7.39
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																		1 (6.7)			1 (1.3)			1 (1.3)	82.00	
肺血栓/肺塞栓 (%)																1 (5.3)					1 (1.3)			1 (1.3)	77.00	
災害・事故死 (%)													1 (9.1)								1 (1.3)			1 (1.3)	63.00	
その他 (%)										1 (100.0)	1 (100.0)		2 (18.2)	2 (15.4)	2 (16.7)	3 (15.8)	2 (13.3)	1 (50.0)			14 (17.7)			14 (17.7)	70.14	11.38
不明 (%)												1 (33.3)		1 (7.7)	2 (16.7)		2 (13.3)			6 (7.6)			6 (7.6)	71.67	10.37	
合計 (%)								1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	3 (100.0)	11 (100.0)	13 (100.0)	12 (100.0)	19 (100.0)	15 (100.0)	2 (100.0)			79 (100.0)			79 (100.0)	71.25	9.81
記載なし																										
総計								1	1	1	1	3	11	13	12	19	15	2			79			79	71.25	9.81

数値下のかっこ内は列方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査