

表 313 2014年死亡患者死亡原因分類(年齢別, 原疾患別)原疾患＝痛風腎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)												1 (33.3)			1 (10.0)	4 (30.8)	5 (31.3)	3 (37.5)	2 (40.0)		16 (23.2)			16 (23.2)	80.69	8.58
脳血管障害 (%)												1 (33.3)	1 (33.3)	3 (27.3)		1 (7.7)	3 (18.8)	1 (12.5)			10 (14.5)			10 (14.5)	72.60	10.04
感染症 (%)													1 (33.3)	1 (9.1)	3 (30.0)	2 (15.4)		2 (25.0)			9 (13.0)			9 (13.0)	75.00	8.43
消化管出血 (%)																	1 (6.3)				1 (1.4)			1 (1.4)	80.00	
悪性腫瘍 (%)														2 (18.2)	2 (20.0)	3 (23.1)	2 (12.5)	1 (12.5)			10 (14.5)			10 (14.5)	76.10	7.02
悪液質/尿毒症 (%)														1 (9.1)	1 (10.0)	2 (15.4)			1 (20.0)		5 (7.2)			5 (7.2)	77.00	9.14
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																	2 (12.5)				2 (2.9)			2 (2.9)	81.50	0.71
肝硬変症 (%)												1 (33.3)									1 (1.4)			1 (1.4)	59.00	
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)														1 (9.1)							1 (1.4)			1 (1.4)	65.00	
災害・事故死 (%)																										
その他 (%)													1 (33.3)	2 (18.2)	1 (10.0)	1 (7.7)	2 (12.5)	1 (12.5)			8 (11.6)			8 (11.6)	74.25	8.96
不明 (%)														1 (9.1)	2 (20.0)		1 (6.3)		2 (40.0)		6 (8.7)			6 (8.7)	79.50	10.01
合計 (%)												3 (100.0)	3 (100.0)	11 (100.0)	10 (100.0)	13 (100.0)	16 (100.0)	8 (100.0)	5 (100.0)		69 (100.0)			69 (100.0)	76.46	8.97
記載なし																										
総計												3	3	11	10	13	16	8	5		69			69	76.46	8.97

数値下のかっこ内は列方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査