

表 316 2014年死亡患者死亡原因分類(年齢別、原疾患別)原疾患=腎・尿路結石

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差		
心不全 (%)														1 (9.1)	3 (33.3)	5 (33.3)	3 (23.1)	1 (33.3)			13 (21.7)			13 (21.7)	81.08	6.51		
脳血管障害 (%)										1 (50.0)				1 (9.1)	1 (20.0)		1 (6.7)				4 (6.7)			4 (6.7)	67.25	15.44		
感染症 (%)														2 (40.0)		6 (40.0)	6 (46.2)				14 (23.3)			14 (23.3)	82.21	5.04		
消化管出血 (%)															1 (6.7)						1 (1.7)			1 (1.7)	83.00			
悪性腫瘍 (%)														3 (27.3)	1 (20.0)			1 (7.7)			6 (10.0)			6 (10.0)	68.83	12.95		
悪液質/尿毒症 (%)														1 (9.1)	2 (22.2)	1 (6.7)				4 (6.7)			4 (6.7)	77.25	6.24			
心筋梗塞 (%)															1 (11.1)					1 (1.7)			1 (1.7)	78.00				
カリウム中毒/頓死 (%)																												
肝硬変症 (%)																					1 (1.7)			1 (1.7)	84.00			
自殺/拒否 (%)														1 (9.1)							1 (1.7)			2 (3.3)	79.50	14.85		
腸閉塞 (%)																												
肺血栓/肺塞栓 (%)																												
災害・事故死 (%)																												
その他 (%)															2 (18.2)	1 (11.1)				2 (15.4)			5 (8.3)			5 (8.3)	76.60	9.56
不明 (%)														2 (100.0)	2 (18.2)	1 (20.0)	2 (22.2)		1 (7.7)	1 (33.3)		9 (15.0)			9 (15.0)	73.67	9.99	
合計 (%)										2 (100.0)		2 (100.0)	5 (100.0)	9 (100.0)	15 (100.0)	13 (100.0)	3 (100.0)		60 (100.0)			60 (100.0)	77.43	9.62				
記載なし																												
総計										2			2	11	5	9	15	13	3		60			60	77.43	9.62		

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査