

表 319 2014年死亡患者死亡原因分類(年齢別、原疾患別)原疾患=骨髄腫

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差			
心不全 (%)										1 (16.7)				2 (11.1)		1 (7.7)				4 (3.6)			4 (3.6)	76.00	15.64				
脳血管障害 (%)										2 (15.4)				1 (3.8)		1 (7.7)				4 (3.6)			4 (3.6)	72.25	11.73				
感染症 (%)										1 (7.7)				2 (12.5)		3 (11.5)		2 (11.1)		1 (7.7)		11 (33.3)			11 (9.9)	72.82	19.32		
消化管出血 (%)										5 (83.3)				7 (53.8)		9 (69.2)		10 (62.5)		16 (61.5)		14 (77.8)		7 (53.8)			70 (63.1)	74.01	9.65
悪性腫瘍 (%)										1 (7.7)				1 (7.7)							1 (0.9)			1 (0.9)	62.00				
悪液質/尿毒症 (%)																					1 (0.9)			1 (0.9)					
心筋梗塞 (%)																					1 (0.9)			1 (0.9)	75.00				
カリウム中毒/頓死 (%)																					1 (0.9)			1 (0.9)					
肝硬変症 (%)																													
自殺/拒否 (%)																													
腸閉塞 (%)																													
肺血栓/肺塞栓 (%)																													
災害・事故死 (%)																													
その他 (%)																					6 (5.4)			6 (5.4)	76.33	7.37			
不明 (%)										1 (100.0)				1 (100.0)		1 (7.7)		1 (6.3)		3 (11.5)		1 (7.7)		6 (15.4)			6 (12.6)	69.43	11.41
合計 (%)					1 (100.0)					1 (100.0)		6 (100.0)	1 (100.0)	13 (100.0)	16 (100.0)	26 (100.0)	18 (100.0)	13 (100.0)	3 (100.0)	111 (100.0)				111 (100.0)	73.35	11.18			
記載なし																													
総計					1				1		6	1	13	13	16	26	18	13	3	111			111	73.35	11.18				

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査