

表 320 2014年死亡患者死亡原因分類(年齢別、原疾患別)原疾患=腎形成不全

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)										1 (100.0)						1 (100.0)					2			2	63.50	26.16	
脳血管障害 (%)								2 (100.0)			1 (33.3)										(11.8)			(11.8)			
感染症 (%)											1 (33.3)										3			3	43.33	8.39	
消化管出血 (%)												1 (100.0)									(17.6)			(17.6)			
悪性腫瘍 (%)													1 (100.0)								5			5	71.20	14.91	
悪液質/尿毒症 (%)																					(29.4)			(29.4)			
カリウム中毒/頓死 (%)																					2			2	78.50	13.44	
心筋梗塞 (%)																					(11.8)			(11.8)			
肝硬変症 (%)																					1			1	25.00		
自殺/拒否 (%)																					(5.9)			(5.9)			
腸閉塞 (%)		1 (100.0)																			1 (5.9)			1 (5.9)		9.00	
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																					3 (17.6)			3 (17.6)		52.67	
不明 (%)																											22.03
合計 (%)	1 (100.0)					1 (100.0)	1 (100.0)	2 (100.0)		1 (100.0)	3 (100.0)		1 (100.0)	1 (100.0)	2 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	17 (100.0)			17 (100.0)		56.59
記載なし																											
総計	1					1	1	2		1	3		1	1	2	1	1	1	1	1	17			17	56.59	23.48	

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査