

表 324 2014年死亡患者死亡原因分類(治療方法別)

死因	施設血液透析	血液透析濾過	血液濾過	血液吸着透析	在宅血液透析	腹膜透析	離脱	生体腎移植	献腎移植	合計	記載なし	総計
心不全	4,410	437	8	15		87				4,957	2,818	7,775
(%)	(26.4)	(31.6)	(42.1)	(20.5)		(23.6)				(26.7)	(25.7)	(26.3)
脳血管障害	1,235	114	2	9		22				1,382	716	2,098
(%)	(7.4)	(8.2)	(10.5)	(12.3)		(6.0)				(7.5)	(6.5)	(7.1)
感染症	3,362	231	2	17	2	89				3,703	2,459	6,162
(%)	(20.1)	(16.7)	(10.5)	(23.3)	(50.0)	(24.1)				(20.0)	(22.4)	(20.9)
消化管出血	252	20		3		3				278	193	471
(%)	(1.5)	(1.4)		(4.1)		(0.8)				(1.5)	(1.8)	(1.6)
悪性腫瘍	1,459	97	1	6	1	30				1,594	1,075	2,669
(%)	(8.7)	(7.0)	(5.3)	(8.2)	(25.0)	(8.1)				(8.6)	(9.8)	(9.0)
悪液質/尿毒症	650	38		1		17				706	464	1,170
(%)	(3.9)	(2.7)		(1.4)		(4.6)				(3.8)	(4.2)	(4.0)
心筋梗塞	703	75		3		19				800	458	1,258
(%)	(4.2)	(5.4)		(4.1)		(5.1)				(4.3)	(4.2)	(4.3)
カリウム中毒/頓死	480	42	1	4		11				538	256	794
(%)	(2.9)	(3.0)	(5.3)	(5.5)		(3.0)				(2.9)	(2.3)	(2.7)
肝硬変症	143	17	1	2		3				166	112	278
(%)	(0.9)	(1.2)	(5.3)	(2.7)		(0.8)				(0.9)	(1.0)	(0.9)
自殺/拒否	127	9		1		1				138	77	215
(%)	(0.8)	(0.7)		(1.4)		(0.3)				(0.7)	(0.7)	(0.7)
腸閉塞	163	8	1			7				179	131	310
(%)	(1.0)	(0.6)	(5.3)			(1.9)				(1.0)	(1.2)	(1.0)
肺血栓/肺塞栓	39			1		1				41	38	79
(%)	(0.2)			(1.4)		(0.3)				(0.2)	(0.3)	(0.3)
災害・事故死	87	9				1				97	60	157
(%)	(0.5)	(0.7)				(0.3)				(0.5)	(0.5)	(0.5)
その他	1,611	101	2	4		35				1,753	997	2,750
(%)	(9.6)	(7.3)	(10.5)	(5.5)		(9.5)				(9.5)	(9.1)	(9.3)
不明	1,978	186	1	7	1	43				2,216	1,123	3,339
(%)	(11.8)	(13.4)	(5.3)	(9.6)	(25.0)	(11.7)				(11.9)	(10.2)	(11.3)
合計	16,699	1,384	19	73	4	369				18,548	10,977	29,525
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)				(100.0)	(100.0)	(100.0)
記載なし												
総計	16,699	1,384	19	73	4	369				18,548	10,977	29,525

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査