

表 328 2014年死亡患者死亡原因の詳細(原疾患との関係)

死因	慢性 糸球体腎炎	慢性 腎盂腎炎	急速進行 性糸球体腎炎	妊娠腎/妊娠 中毒症	その他の分類 不能の腎炎	多発性嚢胞腎	腎硬化症	悪性高血圧	糖尿病性腎症	SLE腎炎	アミロイド腎	痛風腎
心不全	1,761	38	51	19	9	109	687	31	2,696	42	18	10
(%)	(27.4)	(0.6)	(0.8)	(0.3)	(0.1)	(1.7)	(10.7)	(0.5)	(41.9)	(0.7)	(0.3)	(0.2)
肺水腫(溢水)	70		6	1		7	30	2	141	2		3
(%)	(23.8)		(2.0)	(0.3)		(2.4)	(10.2)	(0.7)	(48.0)	(0.7)		(1.0)
急性心筋梗塞(発症30日以内死亡)	148	8	4	3	3	11	92	5	540	7		
(%)	(16.1)	(0.9)	(0.4)	(0.3)	(0.3)	(1.2)	(10.0)	(0.5)	(58.7)	(0.8)		
虚血性心疾患(急性心筋梗塞以外)	68	5	1	1	1	4	41	1	178	2	1	
(%)	(20.1)	(1.5)	(0.3)	(0.3)	(0.3)	(1.2)	(12.1)	(0.3)	(52.7)	(0.6)	(0.3)	
不整脈、伝導障害	83	2	2	3	1	5	44	3	186	2	1	2
(%)	(21.3)	(0.5)	(0.5)	(0.8)	(0.3)	(1.3)	(11.3)	(0.8)	(47.8)	(0.5)	(0.3)	(0.5)
心内膜炎および弁膜症	54	1	3	1	2	1	24		52	3	1	
(%)	(29.3)	(0.5)	(1.6)	(0.5)	(1.1)	(0.5)	(13.0)		(28.3)	(1.6)	(0.5)	
その他の心疾患	121	4	4	1	1	6	63	1	168	5	4	1
(%)	(25.7)	(0.9)	(0.9)	(0.2)	(0.2)	(1.3)	(13.4)	(0.2)	(35.7)	(1.1)	(0.9)	(0.2)
くも膜下出血	44	4	1		1	8	13	1	43	1		1
(%)	(31.2)	(2.8)	(0.7)		(0.7)	(5.7)	(9.2)	(0.7)	(30.5)	(0.7)		(0.7)
脳内出血	254	10	10	2	4	28	92	9	444	2	1	5
(%)	(25.6)	(1.0)	(1.0)	(0.2)	(0.4)	(2.8)	(9.3)	(0.9)	(44.8)	(0.2)	(0.1)	(0.5)
脳梗塞	164	4	11	1	2	16	74	5	331	4		2
(%)	(22.3)	(0.5)	(1.5)	(0.1)	(0.3)	(2.2)	(10.0)	(0.7)	(44.9)	(0.5)		(0.3)
その他の脳血管疾患	60	3	1	1		4	18	1	89			2
(%)	(26.2)	(1.3)	(0.4)	(0.4)		(1.7)	(7.9)	(0.4)	(38.9)			(0.9)
敗血症	531	18	36	8	4	58	237	9	989	35	7	2
(%)	(22.3)	(0.8)	(1.5)	(0.3)	(0.2)	(2.4)	(10.0)	(0.4)	(41.6)	(1.5)	(0.3)	(0.1)
中枢神経系感染症			2			1			6			
(%)			(15.4)			(7.7)			(46.2)			
肺炎	742	31	68	6	13	60	390	21	1,276	17	7	6
(%)	(23.2)	(1.0)	(2.1)	(0.2)	(0.4)	(1.9)	(12.2)	(0.7)	(39.9)	(0.5)	(0.2)	(0.2)
インフルエンザ	1						1		11			
(%)	(6.3)						(6.3)		(68.8)			
尿路感染症	3	1	1			4	7	1	9	1	1	
(%)	(9.7)	(3.2)	(3.2)			(12.9)	(22.6)	(3.2)	(29.0)	(3.2)	(3.2)	
消化管・胆道系感染症・腹膜炎	72	1	1		1	9	26		67	3	1	
(%)	(33.8)	(0.5)	(0.5)		(0.5)	(4.2)	(12.2)		(31.5)	(1.4)	(0.5)	
劇症(急性)ウイルス肝炎	1								4			
(%)	(9.1)								(36.4)			
結核	6						3		16			
(%)	(14.0)						(7.0)		(37.2)			
ヒト免疫不全ウイルス[HIV]感染症									1			
(%)									(100.0)			
その他の感染症	58	2	2	2		8	25	1	113	2	1	1
(%)	(22.8)	(0.8)	(0.8)	(0.8)		(3.1)	(9.8)	(0.4)	(44.5)	(0.8)	(0.4)	(0.4)
中枢神経系の悪性新生物	8						4		7			
(%)	(30.8)						(15.4)		(26.9)			
呼吸器系の悪性新生物	111	4	6		1	8	70	2	181	3		3
(%)	(23.9)	(0.9)	(1.3)		(0.2)	(1.7)	(15.1)	(0.4)	(38.9)	(0.6)		(0.6)
肝癌	85	3	3		1	5	44	7	209			
(%)	(19.3)	(0.7)	(0.7)		(0.2)	(1.1)	(10.0)	(1.6)	(47.5)			
肝癌以外の消化器系の悪性新生物	147	7	3	1	2	8	85	3	274	2		3
(%)	(22.3)	(1.1)	(0.5)	(0.2)	(0.3)	(1.2)	(12.9)	(0.5)	(41.6)	(0.3)		(0.5)
乳房の悪性新生物	16			2		1	6	1	19	2		
(%)	(28.6)			(3.6)		(1.8)	(10.7)	(1.8)	(33.9)	(3.6)		
性器の悪性新生物	26	1	3			1	7	2	35			1
(%)	(22.2)	(0.9)	(2.6)			(0.9)	(6.0)	(1.7)	(29.9)			(0.9)

表 328 2014年死亡患者死亡原因の詳細(原疾患との関係)

死因	慢性 糸球体腎炎	慢性 腎盂腎炎	急速進行 性糸球体腎炎	妊娠腎/妊娠 中毒症	その他の分類 不能の腎炎	多発性嚢胞腎	腎硬化症	悪性高血圧	糖尿病性腎症	SLE腎炎	アミロイド腎	痛風腎
腎の悪性新生物	63			1	2	4	14		32	1		
(%)	(34.1)			(0.5)	(1.1)	(2.2)	(7.6)		(17.3)	(0.5)		
内分泌腺の悪性新生物	14	1				1	5		22			1
(%)	(23.7)	(1.7)				(1.7)	(8.5)		(37.3)			(1.7)
造血・リンパ組織の悪性新生物	52	5	3		1	5	18	1	47	1	2	
(%)	(21.2)	(2.0)	(1.2)		(0.4)	(2.0)	(7.3)	(0.4)	(19.2)	(0.4)	(0.8)	
その他の悪性新生物	90	3	6			8	60	2	133	4	1	2
(%)	(21.5)	(0.7)	(1.4)			(1.9)	(14.4)	(0.5)	(31.8)	(1.0)	(0.2)	(0.5)
ウイルス性肝硬変	39		2			1	9	1	38			
(%)	(34.5)		(1.8)			(0.9)	(8.0)	(0.9)	(33.6)			
ウイルス性以外の肝硬変	44	1	2			5	15	1	65			1
(%)	(26.7)	(0.6)	(1.2)			(3.0)	(9.1)	(0.6)	(39.4)			(0.6)
腸の血行障害	43	1	5			3	19	1	58	2		
(%)	(26.7)	(0.6)	(3.1)			(1.9)	(11.8)	(0.6)	(36.0)	(1.2)		
イレウス	37	1	4			2	12		47	1	1	
(%)	(27.2)	(0.7)	(2.9)			(1.5)	(8.8)		(34.6)	(0.7)	(0.7)	
消化管出血	105	4	8		4	15	48	2	169	7	3	1
(%)	(22.3)	(0.8)	(1.7)		(0.8)	(3.2)	(10.2)	(0.4)	(35.9)	(1.5)	(0.6)	(0.2)
被嚢性腹膜硬化症	7								2			
(%)	(53.8)								(15.4)			
その他の消化器疾患	62	6	5		3	15	31	2	69	2	3	2
(%)	(25.1)	(2.4)	(2.0)		(1.2)	(6.1)	(12.6)	(0.8)	(27.9)	(0.8)	(1.2)	(0.8)
肺梗塞、肺塞栓症	19	1				2	9	3	30		1	1
(%)	(24.1)	(1.3)				(2.5)	(11.4)	(3.8)	(38.0)		(1.3)	(1.3)
悪液質	156	2	7		1	12	85	6	268	3	1	4
(%)	(24.4)	(0.3)	(1.1)		(0.2)	(1.9)	(13.3)	(0.9)	(41.9)	(0.5)	(0.2)	(0.6)
尿毒症	86	2	5		2	6	30	3	130	4		
(%)	(25.9)	(0.6)	(1.5)		(0.6)	(1.8)	(9.0)	(0.9)	(39.2)	(1.2)		
認知症	17						14		20	1		
(%)	(27.0)						(22.2)		(31.7)	(1.6)		
その他の悪液質/尿毒症	35	1	5			3	17		46	3		1
(%)	(25.7)	(0.7)	(3.7)			(2.2)	(12.5)		(33.8)	(2.2)		(0.7)
高カリウム血症	26		2		1	2	11		67	2		1
(%)	(18.6)		(1.4)		(0.7)	(1.4)	(7.9)		(47.9)	(1.4)		(0.7)
原因不明の突然死	125	4	9	3	2	8	71	7	333	5	6	1
(%)	(19.1)	(0.6)	(1.4)	(0.5)	(0.3)	(1.2)	(10.9)	(1.1)	(50.9)	(0.8)	(0.9)	(0.2)
自殺	17		1		1	3	10		42	1		
(%)	(20.5)		(1.2)		(1.2)	(3.6)	(12.0)		(50.6)	(1.2)		
治療拒否(透析拒否)	36	1	4		1	1	14		51	2		
(%)	(27.3)	(0.8)	(3.0)		(0.8)	(0.8)	(10.6)		(38.6)	(1.5)		
災害・事故死	42	2	2			1	22	2	58		1	
(%)	(26.8)	(1.3)	(1.3)			(0.6)	(14.0)	(1.3)	(36.9)		(0.6)	
その他	604	28	34	4	9	56	310	18	861	24	11	6
(%)	(24.1)	(1.1)	(1.4)	(0.2)	(0.4)	(2.2)	(12.4)	(0.7)	(34.4)	(1.0)	(0.4)	(0.2)
合計	6,353	210	323	60	73	515	2,907	155	10,673	198	73	63
(%)	(24.3)	(0.8)	(1.2)	(0.2)	(0.3)	(2.0)	(11.1)	(0.6)	(40.8)	(0.8)	(0.3)	(0.2)
記載なし	789	33	17	8	9	66	285	30	1,356	23	6	6
総計	7,142	243	340	68	82	581	3,192	185	12,029	221	79	69

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査

表 328 2014年死亡患者死亡原因の詳細(原疾患との関係)

死因	先天性代謝異常に基づく腎不全	腎・尿路結核	腎・尿路結石	腎・尿路腫瘍	閉塞性尿路障害	骨髄腫	腎形成不全	不明	再導入	その他	合計	記載なし	総計
心不全	1	4	9	19	16	4	1	736	18	159	6,438		6,438
(%)	(0.0)	(0.1)	(0.1)	(0.3)	(0.2)	(0.1)	(0.0)	(11.4)	(0.3)	(2.5)	(100.0)		
肺水腫(溢水)	1		1	1			1	24		4	294		294
(%)	(0.3)		(0.3)	(0.3)			(0.3)	(8.2)		(1.4)	(100.0)		
急性心筋梗塞(発症30日以内死亡)		1	1	2	4			69	4	18	920		920
(%)		(0.1)	(0.1)	(0.2)	(0.4)			(7.5)	(0.4)	(2.0)	(100.0)		
虚血性心疾患(急性心筋梗塞以外)								23	1	11	338		338
(%)								(6.8)	(0.3)	(3.3)	(100.0)		
不整脈、伝導障害								45	2	8	389		389
(%)								(11.6)	(0.5)	(2.1)	(100.0)		
心内膜炎および弁膜症		2	1	1	2			28		8	184		184
(%)		(1.1)	(0.5)	(0.5)	(1.1)			(15.2)		(4.3)	(100.0)		
その他の心疾患			2	1				63	4	21	470		470
(%)			(0.4)	(0.2)				(13.4)	(0.9)	(4.5)	(100.0)		
くも膜下出血					1	1		19	1	2	141		141
(%)					(0.7)	(0.7)		(13.5)	(0.7)	(1.4)	(100.0)		
脳内出血			2	3	3	2	3	95	5	17	991		991
(%)			(0.2)	(0.3)	(0.3)	(0.2)	(0.3)	(9.6)	(0.5)	(1.7)	(100.0)		
脳梗塞			1	2	1	1		90	1	27	737		737
(%)			(0.1)	(0.3)	(0.1)	(0.1)		(12.2)	(0.1)	(3.7)	(100.0)		
その他の脳血管疾患			1	3	1			35		10	229		229
(%)			(0.4)	(1.3)	(0.4)			(15.3)		(4.4)	(100.0)		
敗血症		1	6	7	4	5	3	300	14	105	2,379		2,379
(%)		(0.0)	(0.3)	(0.3)	(0.2)	(0.2)	(0.1)	(12.6)	(0.6)	(4.4)	(100.0)		
中枢神経系感染症								2		2	13		13
(%)								(15.4)		(15.4)	(100.0)		
肺炎		4	7	8	8	6		432	6	93	3,201		3,201
(%)		(0.1)	(0.2)	(0.2)	(0.2)	(0.2)		(13.5)	(0.2)	(2.9)	(100.0)		
インフルエンザ								3			16		16
(%)								(18.8)			(100.0)		
尿路感染症			1				1			1	31		31
(%)			(3.2)				(3.2)			(3.2)	(100.0)		
消化管・胆道系感染症・腹膜炎	1							25	1	5	213		213
(%)	(0.5)							(11.7)	(0.5)	(2.3)	(100.0)		
劇症(急性)ウイルス肝炎								3		3	11		11
(%)								(27.3)		(27.3)	(100.0)		
結核		1						14		3	43		43
(%)		(2.3)						(32.6)		(7.0)	(100.0)		
ヒト免疫不全ウイルス[HIV]感染症											1		1
(%)											(100.0)		
その他の感染症				2			1	26	1	9	254		254
(%)				(0.8)			(0.4)	(10.2)	(0.4)	(3.5)	(100.0)		
中枢神経系の悪性新生物				1				6			26		26
(%)				(3.8)				(23.1)			(100.0)		
呼吸器系の悪性新生物				5				53	2	16	465		465
(%)				(1.1)				(11.4)	(0.4)	(3.4)	(100.0)		
肝癌		1		4				58	1	19	440		440
(%)		(0.2)		(0.9)				(13.2)	(0.2)	(4.3)	(100.0)		
肝癌以外の消化器系の悪性新生物			2	3	6		1	82	1	28	658		658
(%)			(0.3)	(0.5)	(0.9)		(0.2)	(12.5)	(0.2)	(4.3)	(100.0)		
乳房の悪性新生物								6	1	2	56		56
(%)								(10.7)	(1.8)	(3.6)	(100.0)		
性器の悪性新生物				2	3			24	1	11	117		117
(%)				(1.7)	(2.6)			(20.5)	(0.9)	(9.4)	(100.0)		

表 328 2014年死亡患者死亡原因の詳細(原疾患との関係)

死因	先天性代謝異常に基づく腎不全	腎・尿路結核	腎・尿路結石	腎・尿路腫瘍	閉塞性尿路障害	骨髄腫	腎形成不全	不明	再導入	その他	合計	記載なし	総計
腎の悪性新生物	2		3	31				20	1	11	185		185
(%)	(1.1)		(1.6)	(16.8)				(10.8)	(0.5)	(5.9)	(100.0)		
内分泌腺の悪性新生物				1			1	9	1	3	59		59
(%)				(1.7)			(1.7)	(15.3)	(1.7)	(5.1)	(100.0)		
造血・リンパ組織の悪性新生物					2	67		26		15	245		245
(%)					(0.8)	(27.3)		(10.6)		(6.1)	(100.0)		
その他の悪性新生物		1	1	14	5	3		52		33	418		418
(%)		(0.2)	(0.2)	(3.3)	(1.2)	(0.7)		(12.4)		(7.9)	(100.0)		
ウイルス性肝硬変								10	3	10	113		113
(%)								(8.8)	(2.7)	(8.8)	(100.0)		
ウイルス性以外の肝硬変			1	1			1	21		7	165		165
(%)			(0.6)	(0.6)			(0.6)	(12.7)		(4.2)	(100.0)		
膵の血行障害								23		6	161		161
(%)								(14.3)		(3.7)	(100.0)		
イレウス				2				22	1	6	136		136
(%)				(1.5)				(16.2)	(0.7)	(4.4)	(100.0)		
消化管出血		2	1	1	2			75	1	23	471		471
(%)		(0.4)	(0.2)	(0.2)	(0.4)			(15.9)	(0.2)	(4.9)	(100.0)		
被嚢性腹膜硬化症							1	2		1	13		13
(%)							(7.7)	(15.4)		(7.7)	(100.0)		
その他の消化器疾患						2		29	2	14	247		247
(%)						(0.8)		(11.7)	(0.8)	(5.7)	(100.0)		
肺梗塞、肺塞栓症					1			8		4	79		79
(%)					(1.3)			(10.1)		(5.1)	(100.0)		
悪液質			2	4	1			70	3	14	639		639
(%)			(0.3)	(0.6)	(0.2)			(11.0)	(0.5)	(2.2)	(100.0)		
尿毒症		1	1	1	1			52		8	332		332
(%)		(0.3)	(0.3)	(0.3)	(0.3)			(15.7)		(2.4)	(100.0)		
認知症			1		1			7		2	63		63
(%)			(1.6)		(1.6)			(11.1)		(3.2)	(100.0)		
その他の悪液質/尿毒症				1		1		18		5	136		136
(%)				(0.7)		(0.7)		(13.2)		(3.7)	(100.0)		
高カリウム血症		1		1				19	2	5	140		140
(%)		(0.7)		(0.7)				(13.6)	(1.4)	(3.6)	(100.0)		
原因不明の突然死	2			3	1	1		62	3	8	654		654
(%)	(0.3)			(0.5)	(0.2)	(0.2)		(9.5)	(0.5)	(1.2)	(100.0)		
自殺								5	1	2	83		83
(%)								(6.0)	(1.2)	(2.4)	(100.0)		
治療拒否(透析拒否)			2	1				11	1	7	132		132
(%)			(1.5)	(0.8)				(8.3)	(0.8)	(5.3)	(100.0)		
災害・事故死		1						23		3	157		157
(%)		(0.6)						(14.6)		(1.9)	(100.0)		
その他		4	5	11	6	4	3	371	9	125	2,503		2,503
(%)		(0.2)	(0.2)	(0.4)	(0.2)	(0.2)	(0.1)	(14.8)	(0.4)	(5.0)	(100.0)		
合計	7	24	51	136	69	97	17	3,196	92	894	26,186		26,186
(%)	(0.0)	(0.1)	(0.2)	(0.5)	(0.3)	(0.4)	(0.1)	(12.2)	(0.4)	(3.4)	(100.0)		
記載なし	1	4	9	9	11	14		544	10	109	3,339		3,339
総計	8	28	60	145	80	111	17	3,740	102	1,003	29,525		29,525

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査