

表 330 1983年以降導入2014年末患者原疾患別人数, 平均年齢

原疾患	人 数	記 載 な し	総 計	平 均	標 準 偏 差
慢性糸球体腎炎	93,378	1	93,379	66.61	12.62
(%)	(30.6)	(4.8)	(30.6)		
慢性腎盂腎炎	2,943		2,943	65.52	14.01
(%)	(1.0)		(1.0)		
急速進行性糸球体腎炎	2,475		2,475	68.98	13.00
(%)	(0.8)		(0.8)		
妊娠腎/妊娠中毒症	1,471		1,471	64.17	10.56
(%)	(0.5)		(0.5)		
その他分類不能の腎炎	1,325		1,325	60.39	17.32
(%)	(0.4)		(0.4)		
多発性嚢胞腎	10,944		10,944	65.00	11.31
(%)	(3.6)		(3.6)		
腎硬化症	28,270	1	28,271	74.00	11.77
(%)	(9.3)	(4.8)	(9.3)		
悪性高血圧	2,561		2,561	64.19	14.85
(%)	(0.8)		(0.8)		
糖尿病性腎症	118,047	2	118,049	67.33	11.28
(%)	(38.6)	(9.5)	(38.6)		
SLE腎炎	2,205		2,205	60.74	14.04
(%)	(0.7)		(0.7)		
アミロイド腎	450		450	67.47	11.65
(%)	(0.1)		(0.1)		
痛風腎	1,100	1	1,101	67.56	11.12
(%)	(0.4)	(4.8)	(0.4)		
先天性代謝異常による腎不全	263		263	48.83	17.11
(%)	(0.1)		(0.1)		
腎・尿路結核	188		188	72.70	8.72
(%)	(0.1)		(0.1)		
腎・尿路結石	576		576	70.52	11.10
(%)	(0.2)		(0.2)		
腎・尿路腫瘍	885		885	72.55	10.66
(%)	(0.3)		(0.3)		
閉塞性尿路障害	715	1	716	63.85	16.77
(%)	(0.2)	(4.8)	(0.2)		
骨髄腫	276		276	70.55	12.17
(%)	(0.1)		(0.1)		
腎形成不全	626		626	43.93	19.29
(%)	(0.2)		(0.2)		
不明	27,268	14	27,282	69.84	13.17
(%)	(8.9)	(66.7)	(8.9)		
移植後再導入	2,074		2,074	56.23	12.73
(%)	(0.7)		(0.7)		
その他	7,461	1	7,462	65.66	15.76
(%)	(2.4)	(4.8)	(2.4)		
合計	305,501	21	305,522	67.58	12.55
(%)	(100.0)	(100.0)	(100.0)		
記載なし		1	1	0.00	0.00
総計	305,501	22	305,523	67.58	12.55

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査