

表 493 専従専門医の有無 透析時間(時間/回)別(透析患者全体)

透析時間	なし	あり	合計	記載なし	総計
3.0未満	511	332	843		843
(%)	(60.6)	(39.4)	(100.0)		
3.0～	15,665	14,633	30,298		30,298
(%)	(51.7)	(48.3)	(100.0)		
3.5～	9,930	11,281	21,211		21,211
(%)	(46.8)	(53.2)	(100.0)		
4.0～	80,591	107,199	187,790		187,790
(%)	(42.9)	(57.1)	(100.0)		
4.5～	6,317	9,759	16,076		16,076
(%)	(39.3)	(60.7)	(100.0)		
5.0～	7,261	13,156	20,417		20,417
(%)	(35.6)	(64.4)	(100.0)		
5.5～	297	570	867		867
(%)	(34.3)	(65.7)	(100.0)		
6.0～	370	1,032	1,402		1,402
(%)	(26.4)	(73.6)	(100.0)		
6.5～	22	64	86		86
(%)	(25.6)	(74.4)	(100.0)		
7.0～	36	611	647		647
(%)	(5.6)	(94.4)	(100.0)		
合計	121,000	158,637	279,637		279,637
(%)	(43.3)	(56.7)	(100.0)		
記載なし	12,340	18,130	30,470	1	30,471
(%)	(40.5)	(59.5)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	3.93	4.03	3.98		3.98
標準偏差	0.50	0.55	0.53		0.53

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査