

表 495 専従専門医の有無 身長(cm)別(透析患者全体)

身長	なし	あり	合計	記載なし	総計
130未満	167	186	353	1	354
(%)	(47.3)	(52.7)	(100.0)		
130～	325	371	696		696
(%)	(46.7)	(53.3)	(100.0)		
135～	1,325	1,512	2,837		2,837
(%)	(46.7)	(53.3)	(100.0)		
140～	4,657	5,640	10,297		10,297
(%)	(45.2)	(54.8)	(100.0)		
145～	10,560	13,113	23,673		23,673
(%)	(44.6)	(55.4)	(100.0)		
150～	16,831	21,404	38,235		38,235
(%)	(44.0)	(56.0)	(100.0)		
155～	19,197	25,275	44,472		44,472
(%)	(43.2)	(56.8)	(100.0)		
160～	23,203	30,447	53,650		53,650
(%)	(43.2)	(56.8)	(100.0)		
165～	19,959	28,029	47,988		47,988
(%)	(41.6)	(58.4)	(100.0)		
170～	12,838	18,236	31,074		31,074
(%)	(41.3)	(58.7)	(100.0)		
175～	4,537	6,782	11,319		11,319
(%)	(40.1)	(59.9)	(100.0)		
180～	1,165	1,814	2,979		2,979
(%)	(39.1)	(60.9)	(100.0)		
185～	168	297	465		465
(%)	(36.1)	(63.9)	(100.0)		
190～	33	44	77		77
(%)	(42.9)	(57.1)	(100.0)		
合計	114,965	153,150	268,115	1	268,116
(%)	(42.9)	(57.1)	(100.0)		
記載なし	18,375	23,617	41,992		41,992
(%)	(43.8)	(56.2)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	159.72	160.29	160.05	97.00	160.05
標準偏差	9.65	9.64	9.65		9.65

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査