

表 499 専従専門医の有無 透析前アルブミン濃度 (g/dL) 別 (透析患者全体)

透析前アルブミン 濃度	なし	あり	合計	記載なし	総計
1.5未満	96	89	185		185
(%)	(51.9)	(48.1)	(100.0)		
1.5～	524	465	989		989
(%)	(53.0)	(47.0)	(100.0)		
2.0～	2,083	1,977	4,060	1	4,061
(%)	(51.3)	(48.7)	(100.0)		
2.5～	7,536	7,900	15,436		15,436
(%)	(48.8)	(51.2)	(100.0)		
3.0～	29,767	36,341	66,108		66,108
(%)	(45.0)	(55.0)	(100.0)		
3.5～	56,606	78,629	135,235		135,235
(%)	(41.9)	(58.1)	(100.0)		
4.0～	20,457	30,051	50,508		50,508
(%)	(40.5)	(59.5)	(100.0)		
4.5～	1,177	1,737	2,914		2,914
(%)	(40.4)	(59.6)	(100.0)		
5.0～	89	114	203		203
(%)	(43.8)	(56.2)	(100.0)		
合計	118,335	157,303	275,638	1	275,639
(%)	(42.9)	(57.1)	(100.0)		
記載なし	15,005	19,464	34,469		34,469
(%)	(43.5)	(56.5)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	3.57	3.62	3.60	2.20	3.60
標準偏差	0.46	0.43	0.44		0.44

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査