

表 500 専従専門医の有無 透析前CRP濃度(mg/dL)別(透析患者全体)

透析前CRP濃度	なし	あり	合計	記載なし	総計
0.2未満	57,032	78,276	135,308	1	135,309
(%)	(42.1)	(57.9)	(100.0)		
0.2～	19,086	25,721	44,807		44,807
(%)	(42.6)	(57.4)	(100.0)		
0.5～	9,777	12,980	22,757		22,757
(%)	(43.0)	(57.0)	(100.0)		
1.0～	6,446	8,302	14,748		14,748
(%)	(43.7)	(56.3)	(100.0)		
2.0～	4,026	4,979	9,005		9,005
(%)	(44.7)	(55.3)	(100.0)		
4.0～	1,402	1,842	3,244		3,244
(%)	(43.2)	(56.8)	(100.0)		
6.0～	722	854	1,576		1,576
(%)	(45.8)	(54.2)	(100.0)		
8.0～	371	499	870		870
(%)	(42.6)	(57.4)	(100.0)		
10.0～	401	633	1,034		1,034
(%)	(38.8)	(61.2)	(100.0)		
15.0～	296	397	693		693
(%)	(42.7)	(57.3)	(100.0)		
合計	99,559	134,483	234,042	1	234,043
(%)	(42.5)	(57.5)	(100.0)		
記載なし	33,781	42,284	76,065		76,065
(%)	(44.4)	(55.6)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	0.64	0.63	0.63	0.03	0.63
標準偏差	1.88	1.95	1.92		1.92

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査