

表 501 専従専門医の有無 透析前補正カルシウム濃度 (mg/dL) 別 (透析患者全体)

透析前補正 カルシウム濃度	なし	あり	合計	記載なし	総計
7.0未満	551	591	1,142		1,142
(%)	(48.2)	(51.8)	(100.0)		
7.0～	713	936	1,649		1,649
(%)	(43.2)	(56.8)	(100.0)		
7.5～	2,769	3,630	6,399		6,399
(%)	(43.3)	(56.7)	(100.0)		
8.0～	8,551	11,888	20,439		20,439
(%)	(41.8)	(58.2)	(100.0)		
8.4～	2,429	3,443	5,872		5,872
(%)	(41.4)	(58.6)	(100.0)		
8.5～	27,984	39,335	67,319		67,319
(%)	(41.6)	(58.4)	(100.0)		
9.0～	34,450	47,554	82,004		82,004
(%)	(42.0)	(58.0)	(100.0)		
9.5～	23,455	30,514	53,969		53,969
(%)	(43.5)	(56.5)	(100.0)		
10.0～	2,965	3,633	6,598		6,598
(%)	(44.9)	(55.1)	(100.0)		
10.1～	7,885	9,091	16,976	1	16,977
(%)	(46.4)	(53.6)	(100.0)		
10.5～	4,049	4,221	8,270		8,270
(%)	(49.0)	(51.0)	(100.0)		
11.0～	1,457	1,324	2,781		2,781
(%)	(52.4)	(47.6)	(100.0)		
11.5～	532	441	973		973
(%)	(54.7)	(45.3)	(100.0)		
12.0～	371	310	681		681
(%)	(54.5)	(45.5)	(100.0)		
合計	118,161	156,911	275,072	1	275,073
(%)	(43.0)	(57.0)	(100.0)		
記載なし	15,179	19,856	35,035		35,035
(%)	(43.3)	(56.7)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	9.21	9.17	9.19	10.30	9.19
標準偏差	0.77	0.72	0.74		0.74

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査