

表 502 専従専門医の有無 透析前リン濃度 (mg/dL) 別 (透析患者全体)

透析前リン濃度	なし	あり	合計	記載なし	総計
3未満	5,409	5,477	10,886	1	10,886
(%)	(49.7)	(50.3)	(100.0)		
3.0～	6,163	7,419	13,582		13,582
(%)	(45.4)	(54.6)	(100.0)		
3.5～	10,303	13,379	23,682		23,682
(%)	(43.5)	(56.5)	(100.0)		
4.0～	15,136	20,145	35,281		35,282
(%)	(42.9)	(57.1)	(100.0)		
4.5～	17,656	24,403	42,059		42,059
(%)	(42.0)	(58.0)	(100.0)		
5.0～	17,786	24,932	42,718		42,718
(%)	(41.6)	(58.4)	(100.0)		
5.5～	15,408	21,297	36,705		36,705
(%)	(42.0)	(58.0)	(100.0)		
6.0～	2,605	3,656	6,261		6,261
(%)	(41.6)	(58.4)	(100.0)		
6.1～	9,014	12,095	21,109		21,109
(%)	(42.7)	(57.3)	(100.0)		
6.5～	7,808	10,470	18,278		18,278
(%)	(42.7)	(57.3)	(100.0)		
7.0～	4,890	6,387	11,277		11,277
(%)	(43.4)	(56.6)	(100.0)		
7.5～	3,028	3,830	6,858		6,858
(%)	(44.2)	(55.8)	(100.0)		
8.0～	1,751	2,302	4,053		4,053
(%)	(43.2)	(56.8)	(100.0)		
8.5～	1,157	1,415	2,572		2,572
(%)	(45.0)	(55.0)	(100.0)		
9.0～	1,910	2,167	4,077		4,077
(%)	(46.8)	(53.2)	(100.0)		
合計	120,024	159,374	279,398	1	279,399
(%)	(43.0)	(57.0)	(100.0)		
記載なし	13,316	17,393	30,709		30,709
(%)	(43.4)	(56.6)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	5.20	5.22	5.21	4.00	5.21
標準偏差	1.47	1.39	1.43		1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査