

表 504 専従専門医の有無 PTH値 (pg/mL) 別 (透析患者全体、PTH測定法:whole-PTH)

PTH値	なし	あり	合計	記載なし	総計
20未満	601	1,293	1,894		1,894
(%)	(31.7)	(68.3)	(100.0)		
20～	704	1,627	2,331		2,331
(%)	(30.2)	(69.8)	(100.0)		
35～	588	1,455	2,043		2,043
(%)	(28.8)	(71.2)	(100.0)		
50～	851	2,302	3,153		3,153
(%)	(27.0)	(73.0)	(100.0)		
75～	718	2,030	2,748		2,748
(%)	(26.1)	(73.9)	(100.0)		
100～	557	1,741	2,298		2,298
(%)	(24.2)	(75.8)	(100.0)		
125～	404	1,135	1,539		1,539
(%)	(26.3)	(73.7)	(100.0)		
150～	11	37	48		48
(%)	(22.9)	(77.1)	(100.0)		
151～	339	924	1,263		1,263
(%)	(26.8)	(73.2)	(100.0)		
180～	240	635	875		875
(%)	(27.4)	(72.6)	(100.0)		
210～	182	446	628		628
(%)	(29.0)	(71.0)	(100.0)		
240～	203	471	674		674
(%)	(30.1)	(69.9)	(100.0)		
300～	95	189	284		284
(%)	(33.5)	(66.5)	(100.0)		
350～	59	120	179		179
(%)	(33.0)	(67.0)	(100.0)		
400～	73	129	202		202
(%)	(36.1)	(63.9)	(100.0)		
500～	83	164	247		247
(%)	(33.6)	(66.4)	(100.0)		
合計	5,708	14,698	20,406		20,406
(%)	(28.0)	(72.0)	(100.0)		
記載なし	136	521	657		657
(%)	(20.7)	(79.3)	(100.0)		
総計	5,844	15,219	21,063		21,063
(%)	(27.7)	(72.3)	(100.0)		
平均	109.67	107.36	108.01		108.01
標準偏差	117.04	112.82	114.01		114.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査