

表 505 専従専門医の有無 透析前ヘモグロビン濃度 (g/dL) 別 (透析患者全体)

透析前ヘモグロビン 濃度	なし	あり	合計	記載なし	総計
5未満	107	196	303		303
(%)	(35.3)	(64.7)	(100.0)		
5～	186	243	429		429
(%)	(43.4)	(56.6)	(100.0)		
6～	586	574	1,160		1,160
(%)	(50.5)	(49.5)	(100.0)		
7～	2,180	2,254	4,434	1	4,435
(%)	(49.2)	(50.8)	(100.0)		
8～	7,038	7,516	14,554		14,554
(%)	(48.4)	(51.6)	(100.0)		
9～	20,733	25,256	45,989		45,989
(%)	(45.1)	(54.9)	(100.0)		
10～	39,816	54,096	93,912		93,912
(%)	(42.4)	(57.6)	(100.0)		
11～	32,108	46,065	78,173		78,173
(%)	(41.1)	(58.9)	(100.0)		
12～	12,218	16,636	28,854		28,854
(%)	(42.3)	(57.7)	(100.0)		
13～	3,203	4,105	7,308		7,308
(%)	(43.8)	(56.2)	(100.0)		
14～	1,218	1,455	2,673		2,673
(%)	(45.6)	(54.4)	(100.0)		
合計	119,393	158,396	277,789	1	277,790
(%)	(43.0)	(57.0)	(100.0)		
記載なし	13,947	18,371	32,318		32,318
(%)	(43.2)	(56.8)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	10.67	10.74	10.71	7.30	10.71
標準偏差	1.31	1.26	1.28		1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査