

表 507 専従専門医の有無 HDL-C濃度(mg/dL)別(透析患者全体)

HDL-C濃度	なし	あり	合計	記載なし	総計
20未満	363	432	795	1	796
(%)	(45.7)	(54.3)	(100.0)		
20～	6,769	8,854	15,623		15,623
(%)	(43.3)	(56.7)	(100.0)		
30～	19,801	26,046	45,847		45,847
(%)	(43.2)	(56.8)	(100.0)		
40～	22,283	29,069	51,352		51,352
(%)	(43.4)	(56.6)	(100.0)		
50～	16,226	21,335	37,561		37,561
(%)	(43.2)	(56.8)	(100.0)		
60～	9,305	12,931	22,236		22,236
(%)	(41.8)	(58.2)	(100.0)		
70～	4,605	6,387	10,992		10,992
(%)	(41.9)	(58.1)	(100.0)		
80～	2,238	2,991	5,229		5,229
(%)	(42.8)	(57.2)	(100.0)		
90～	984	1,347	2,331		2,331
(%)	(42.2)	(57.8)	(100.0)		
100～	860	1,141	2,001		2,001
(%)	(43.0)	(57.0)	(100.0)		
合計	83,434	110,533	193,967	1	193,968
(%)	(43.0)	(57.0)	(100.0)		
記載なし	49,906	66,234	116,140		116,140
(%)	(43.0)	(57.0)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	48.61	48.86	48.75	17.00	48.75
標準偏差	16.65	16.74	16.71		16.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査