

表 510 専従専門医の有無 透析前脈拍(拍/分)別(透析患者全体)

透析前脈拍	なし	あり	合計	記載なし	総計
40未満	85	100	185		185
(%)	(45.9)	(54.1)	(100.0)		
40～	1,152	1,229	2,381		2,381
(%)	(48.4)	(51.6)	(100.0)		
50～	8,848	10,664	19,512		19,512
(%)	(45.3)	(54.7)	(100.0)		
60～	26,768	34,013	60,781		60,781
(%)	(44.0)	(56.0)	(100.0)		
70～	31,525	41,747	73,272		73,272
(%)	(43.0)	(57.0)	(100.0)		
80～	19,972	26,842	46,814		46,814
(%)	(42.7)	(57.3)	(100.0)		
90～	8,807	11,783	20,590		20,590
(%)	(42.8)	(57.2)	(100.0)		
100～	2,672	3,721	6,393		6,393
(%)	(41.8)	(58.2)	(100.0)		
110～	626	902	1,528		1,528
(%)	(41.0)	(59.0)	(100.0)		
120～	265	389	654	1	655
(%)	(40.5)	(59.5)	(100.0)		
合計	100,720	131,390	232,110	1	232,111
(%)	(43.4)	(56.6)	(100.0)		
記載なし	32,620	45,377	77,997		77,997
(%)	(41.8)	(58.2)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	74.41	74.90	74.69	126.00	74.69
標準偏差	12.84	12.83	12.84		12.84

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査