

表 516 専従専門医の有無 体重(kg)別(施設血液透析患者)

体重	なし	あり	合計	記載なし	総計
20未満	1	4	5		5
(%)	(20.0)	(80.0)	(100.0)		
20～	196	245	441		441
(%)	(44.4)	(55.6)	(100.0)		
30～	6,524	7,314	13,838		13,838
(%)	(47.1)	(52.9)	(100.0)		
40～	24,404	29,372	53,776		53,776
(%)	(45.4)	(54.6)	(100.0)		
50～	32,207	41,941	74,148		74,148
(%)	(43.4)	(56.6)	(100.0)		
60～	22,131	30,887	53,018		53,018
(%)	(41.7)	(58.3)	(100.0)		
70～	9,468	13,767	23,235		23,235
(%)	(40.7)	(59.3)	(100.0)		
80～	3,345	5,189	8,534		8,534
(%)	(39.2)	(60.8)	(100.0)		
90～	1,250	1,951	3,201		3,201
(%)	(39.1)	(60.9)	(100.0)		
100～	449	665	1,114		1,114
(%)	(40.3)	(59.7)	(100.0)		
110～	182	321	503		503
(%)	(36.2)	(63.8)	(100.0)		
120～	129	211	340		340
(%)	(37.9)	(62.1)	(100.0)		
合計	100,286	131,867	232,153		232,153
(%)	(43.2)	(56.8)	(100.0)		
記載なし	10,206	13,282	23,488		23,488
(%)	(43.5)	(56.5)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	57.19	58.30	57.82		57.82
標準偏差	13.25	13.54	13.42		13.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査