

表 517 専従専門医の有無 透析前BUN(mg/dL)別(施設血液透析患者)

透析前BUN	なし	あり	合計	記載なし	総計
20未満	372	342	714		714
(%)	(52.1)	(47.9)	(100.0)		
20～	1,808	2,025	3,833		3,833
(%)	(47.2)	(52.8)	(100.0)		
30～	6,842	7,919	14,761		14,761
(%)	(46.4)	(53.6)	(100.0)		
40～	16,197	20,279	36,476		36,476
(%)	(44.4)	(55.6)	(100.0)		
50～	24,316	32,456	56,772		56,772
(%)	(42.8)	(57.2)	(100.0)		
60～	23,681	32,500	56,181		56,181
(%)	(42.2)	(57.8)	(100.0)		
70～	15,775	21,222	36,997		36,997
(%)	(42.6)	(57.4)	(100.0)		
80～	7,460	9,894	17,354		17,354
(%)	(43.0)	(57.0)	(100.0)		
90～	2,771	3,647	6,418		6,418
(%)	(43.2)	(56.8)	(100.0)		
100～	844	1,080	1,924		1,924
(%)	(43.9)	(56.1)	(100.0)		
110～	268	324	592		592
(%)	(45.3)	(54.7)	(100.0)		
120～	174	183	357		357
(%)	(48.7)	(51.3)	(100.0)		
合計	100,508	131,871	232,379		232,379
(%)	(43.3)	(56.7)	(100.0)		
記載なし	9,984	13,278	23,262		23,262
(%)	(42.9)	(57.1)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	60.40	60.88	60.68		60.68
標準偏差	16.22	15.74	15.95		15.95

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査