

表 518 専従専門医の有無 透析前クレアチニン濃度 (mg/dL) 別 (施設血液透析患者)

透析前クレアチニン濃度	なし	あり	合計	記載なし	総計
2未満	160	153	313		313
(%)	(51.1)	(48.9)	(100.0)		
2～	2,166	2,210	4,376		4,376
(%)	(49.5)	(50.5)	(100.0)		
4～	7,770	8,449	16,219		16,219
(%)	(47.9)	(52.1)	(100.0)		
6～	17,728	21,250	38,978		38,978
(%)	(45.5)	(54.5)	(100.0)		
8～	26,837	34,946	61,783		61,783
(%)	(43.4)	(56.6)	(100.0)		
10～	24,410	33,580	57,990		57,990
(%)	(42.1)	(57.9)	(100.0)		
12～	14,071	20,128	34,199		34,199
(%)	(41.1)	(58.9)	(100.0)		
14～	5,510	8,088	13,598		13,598
(%)	(40.5)	(59.5)	(100.0)		
16～	1,898	2,857	4,755		4,755
(%)	(39.9)	(60.1)	(100.0)		
合計	100,550	131,661	232,211		232,211
(%)	(43.3)	(56.7)	(100.0)		
記載なし	9,942	13,488	23,430		23,430
(%)	(42.4)	(57.6)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	9.69	9.94	9.83		9.83
標準偏差	2.97	2.92	2.94		2.94

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査