

表 521 専従専門医の有無 透析前補正カルシウム濃度 (mg/dL) 別 (施設血液透析患者)

透析前補正 カルシウム濃度	なし	あり	合計	記載なし	総計
7.0未満	440	491	931		931
(%)	(47.3)	(52.7)	(100.0)		
7.0～	574	757	1,331		1,331
(%)	(43.1)	(56.9)	(100.0)		
7.5～	2,320	2,925	5,245		5,245
(%)	(44.2)	(55.8)	(100.0)		
8.0～	7,196	9,866	17,062		17,062
(%)	(42.2)	(57.8)	(100.0)		
8.4～	2,020	2,878	4,898		4,898
(%)	(41.2)	(58.8)	(100.0)		
8.5～	23,575	32,857	56,432		56,432
(%)	(41.8)	(58.2)	(100.0)		
9.0～	28,780	39,231	68,011		68,011
(%)	(42.3)	(57.7)	(100.0)		
9.5～	19,321	24,765	44,086		44,086
(%)	(43.8)	(56.2)	(100.0)		
10.0～	2,459	2,957	5,416		5,416
(%)	(45.4)	(54.6)	(100.0)		
10.1～	6,449	7,283	13,732		13,732
(%)	(47.0)	(53.0)	(100.0)		
10.5～	3,318	3,374	6,692		6,692
(%)	(49.6)	(50.4)	(100.0)		
11.0～	1,200	1,069	2,269		2,269
(%)	(52.9)	(47.1)	(100.0)		
11.5～	429	373	802		802
(%)	(53.5)	(46.5)	(100.0)		
12.0～	312	263	575		575
(%)	(54.3)	(45.7)	(100.0)		
合計	98,393	129,089	227,482		227,482
(%)	(43.3)	(56.7)	(100.0)		
記載なし	12,099	16,060	28,159		28,159
(%)	(43.0)	(57.0)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	9.21	9.16	9.18		9.18
標準偏差	0.77	0.72	0.74		0.74

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査