

表 522 専従専門医の有無 透析前リン濃度(mg/dL)別(施設血液透析患者)

透析前リン濃度	なし	あり	合計	記載なし	総計
3未満	4,767	4,742	9,509		9,509
(%)	(50.1)	(49.9)	(100.0)		
3.0～	5,299	6,281	11,580		11,580
(%)	(45.8)	(54.2)	(100.0)		
3.5～	8,740	11,275	20,015		20,015
(%)	(43.7)	(56.3)	(100.0)		
4.0～	12,807	16,736	29,543		29,543
(%)	(43.4)	(56.6)	(100.0)		
4.5～	14,764	20,141	34,905		34,905
(%)	(42.3)	(57.7)	(100.0)		
5.0～	14,707	20,444	35,151		35,151
(%)	(41.8)	(58.2)	(100.0)		
5.5～	12,722	17,419	30,141		30,141
(%)	(42.2)	(57.8)	(100.0)		
6.0～	2,153	2,956	5,109		5,109
(%)	(42.1)	(57.9)	(100.0)		
6.1～	7,430	9,726	17,156		17,156
(%)	(43.3)	(56.7)	(100.0)		
6.5～	6,395	8,510	14,905		14,905
(%)	(42.9)	(57.1)	(100.0)		
7.0～	4,029	5,170	9,199		9,199
(%)	(43.8)	(56.2)	(100.0)		
7.5～	2,466	3,073	5,539		5,539
(%)	(44.5)	(55.5)	(100.0)		
8.0～	1,437	1,861	3,298		3,298
(%)	(43.6)	(56.4)	(100.0)		
8.5～	940	1,144	2,084		2,084
(%)	(45.1)	(54.9)	(100.0)		
9.0～	1,530	1,746	3,276		3,276
(%)	(46.7)	(53.3)	(100.0)		
合計	100,186	131,224	231,410		231,410
(%)	(43.3)	(56.7)	(100.0)		
記載なし	10,306	13,925	24,231		24,231
(%)	(42.5)	(57.5)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	5.17	5.20	5.19		5.19
標準偏差	1.47	1.40	1.43		1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査