

表 523 専従専門医の有無 PTH値(pg/mL)別(施設血液透析患者、PTH測定法:intact-PTH)

PTH値	なし	あり	合計	記載なし	総計
20未満	4,946	4,926	9,872		9,872
(%)	(50.1)	(49.9)	(100.0)		
20～	6,848	7,422	14,270		14,270
(%)	(48.0)	(52.0)	(100.0)		
40～	6,946	8,170	15,116		15,116
(%)	(46.0)	(54.0)	(100.0)		
60～	7,280	8,833	16,113		16,113
(%)	(45.2)	(54.8)	(100.0)		
80～	7,193	9,068	16,261		16,261
(%)	(44.2)	(55.8)	(100.0)		
100～	7,003	8,851	15,854		15,854
(%)	(44.2)	(55.8)	(100.0)		
120～	6,646	8,501	15,147		15,147
(%)	(43.9)	(56.1)	(100.0)		
140～	6,061	7,974	14,035		14,035
(%)	(43.2)	(56.8)	(100.0)		
160～	5,469	7,129	12,598		12,598
(%)	(43.4)	(56.6)	(100.0)		
180～	12,363	16,143	28,506		28,506
(%)	(43.4)	(56.6)	(100.0)		
240～	162	189	351		351
(%)	(46.2)	(53.8)	(100.0)		
241～	7,031	9,223	16,254		16,254
(%)	(43.3)	(56.7)	(100.0)		
300～	4,091	5,275	9,366		9,366
(%)	(43.7)	(56.3)	(100.0)		
360～	2,372	2,997	5,369		5,369
(%)	(44.2)	(55.8)	(100.0)		
420～	1,379	1,776	3,155		3,155
(%)	(43.7)	(56.3)	(100.0)		
480～	329	441	770		770
(%)	(42.7)	(57.3)	(100.0)		
500～	573	685	1,258		1,258
(%)	(45.5)	(54.5)	(100.0)		
540～	616	744	1,360		1,360
(%)	(45.3)	(54.7)	(100.0)		
600～	1,000	1,228	2,228		2,228
(%)	(44.9)	(55.1)	(100.0)		
800～	781	903	1,684		1,684
(%)	(46.4)	(53.6)	(100.0)		
合計	89,089	110,478	199,567		199,567
(%)	(44.6)	(55.4)	(100.0)		
記載なし	2,285	2,733	5,018		5,018
(%)	(45.5)	(54.5)	(100.0)		
総計	91,374	113,211	204,585		204,585
(%)	(44.7)	(55.3)	(100.0)		
平均	168.12	171.82	170.17		170.17
標準偏差	165.27	157.54	161.05		161.05

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査