

表 525 専従専門医の有無 透析前ヘモグロビン濃度 (g/dL) 別 (施設血液透析患者)

透析前ヘモグロビン 濃度	なし	あり	合計	記載なし	総計
5未満	90	179	269		269
(%)	(33.5)	(66.5)	(100.0)		
5～	172	208	380		380
(%)	(45.3)	(54.7)	(100.0)		
6～	512	499	1,011		1,011
(%)	(50.6)	(49.4)	(100.0)		
7～	1,890	1,932	3,822		3,822
(%)	(49.5)	(50.5)	(100.0)		
8～	6,054	6,356	12,410		12,410
(%)	(48.8)	(51.2)	(100.0)		
9～	17,598	21,112	38,710		38,710
(%)	(45.5)	(54.5)	(100.0)		
10～	33,353	44,908	78,261		78,261
(%)	(42.6)	(57.4)	(100.0)		
11～	26,525	37,511	64,036		64,036
(%)	(41.4)	(58.6)	(100.0)		
12～	9,904	13,418	23,322		23,322
(%)	(42.5)	(57.5)	(100.0)		
13～	2,583	3,207	5,790		5,790
(%)	(44.6)	(55.4)	(100.0)		
14～	948	1,066	2,014		2,014
(%)	(47.1)	(52.9)	(100.0)		
合計	99,629	130,396	230,025		230,025
(%)	(43.3)	(56.7)	(100.0)		
記載なし	10,863	14,753	25,616		25,616
(%)	(42.4)	(57.6)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	10.64	10.72	10.68		10.68
標準偏差	1.31	1.26	1.28		1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査