

表 527 専従専門医の有無 HDL-C濃度(mg/dL)別(施設血液透析患者)

HDL-C濃度	なし	あり	合計	記載なし	総計
20未満	324	377	701		701
(%)	(46.2)	(53.8)	(100.0)		
20～	5,805	7,521	13,326		13,326
(%)	(43.6)	(56.4)	(100.0)		
30～	16,683	21,708	38,391		38,391
(%)	(43.5)	(56.5)	(100.0)		
40～	18,489	23,920	42,409		42,409
(%)	(43.6)	(56.4)	(100.0)		
50～	13,179	17,143	30,322		30,322
(%)	(43.5)	(56.5)	(100.0)		
60～	7,391	10,151	17,542		17,542
(%)	(42.1)	(57.9)	(100.0)		
70～	3,630	4,913	8,543		8,543
(%)	(42.5)	(57.5)	(100.0)		
80～	1,765	2,236	4,001		4,001
(%)	(44.1)	(55.9)	(100.0)		
90～	766	1,035	1,801		1,801
(%)	(42.5)	(57.5)	(100.0)		
100～	675	854	1,529		1,529
(%)	(44.1)	(55.9)	(100.0)		
合計	68,707	89,858	158,565		158,565
(%)	(43.3)	(56.7)	(100.0)		
記載なし	41,785	55,291	97,076		97,076
(%)	(43.0)	(57.0)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	48.19	48.34	48.28		48.28
標準偏差	16.56	16.53	16.54		16.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査