

表 528 専従専門医の有無 透析前収縮期血圧(mmHg)別(施設血液透析患者)

透析前収縮 期血圧	なし	あり	合計	記載なし	総計
60未満	14	22	36		36
(%)	(38.9)	(61.1)	(100.0)		
60～	119	172	291		291
(%)	(40.9)	(59.1)	(100.0)		
80～	1,022	1,481	2,503		2,503
(%)	(40.8)	(59.2)	(100.0)		
100～	5,926	7,901	13,827		13,827
(%)	(42.9)	(57.1)	(100.0)		
120～	18,282	25,186	43,468		43,468
(%)	(42.1)	(57.9)	(100.0)		
140～	28,354	38,560	66,914		66,914
(%)	(42.4)	(57.6)	(100.0)		
160～	21,188	27,450	48,638		48,638
(%)	(43.6)	(56.4)	(100.0)		
180～	8,338	10,345	18,683		18,683
(%)	(44.6)	(55.4)	(100.0)		
200～	2,102	2,568	4,670		4,670
(%)	(45.0)	(55.0)	(100.0)		
220～	424	506	930		930
(%)	(45.6)	(54.4)	(100.0)		
合計	85,769	114,191	199,960		199,960
(%)	(42.9)	(57.1)	(100.0)		
記載なし	24,723	30,958	55,681		55,681
(%)	(44.4)	(55.6)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	151.96	151.24	151.55		151.55
標準偏差	24.20	23.90	24.03		24.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査