

表 531 専従専門医の有無 nPCR(g/kg/day)別(施設血液透析患者)

nPCR	なし	あり	合計	記載なし	総計
0.4未満	344	308	652		652
(%)	(52.8)	(47.2)	(100.0)		
0.4～	6,898	7,347	14,245		14,245
(%)	(48.4)	(51.6)	(100.0)		
0.6～	32,101	39,291	71,392		71,392
(%)	(45.0)	(55.0)	(100.0)		
0.8～	38,109	51,387	89,496		89,496
(%)	(42.6)	(57.4)	(100.0)		
1.0～	14,487	19,789	34,276		34,276
(%)	(42.3)	(57.7)	(100.0)		
1.2～	2,475	3,372	5,847		5,847
(%)	(42.3)	(57.7)	(100.0)		
1.4～	287	374	661		661
(%)	(43.4)	(56.6)	(100.0)		
1.6～	55	42	97		97
(%)	(56.7)	(43.3)	(100.0)		
1.8～	12	26	38		38
(%)	(31.6)	(68.4)	(100.0)		
2.0～	23	29	52		52
(%)	(44.2)	(55.8)	(100.0)		
合計	94,791	121,965	216,756		216,756
(%)	(43.7)	(56.3)	(100.0)		
記載なし	15,701	23,184	38,885		38,885
(%)	(40.4)	(59.6)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	0.84	0.85	0.85		0.85
標準偏差	0.18	0.18	0.18		0.18

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査